

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12N 15/53, 15/82, 15/81, 9/06, 1/21, 1/19, 5/10, A01H 5/00, C12Q 1/26</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/06571</b> <b>(43) International Publication Date:</b> 11 February 1999 (11.02.99)
<b>(21) International Application Number:</b> PCT/US98/15844 <b>(22) International Filing Date:</b> 30 July 1998 (30.07.98)  <b>(30) Priority Data:</b> 60/054,268      30 July 1997 (30.07.97)      US 09/124,541      29 July 1998 (29.07.98)      US  <b>(71) Applicant:</b> THE CURATORS OF THE UNIVERSITY OF MISSOURI [US/US]; 509 Lewis Hall, Columbia, MO 65211 (US).  <b>(72) Inventor:</b> MORRIS, Roy, O.; 4938 Bethel Street, Columbia, MO 65203 (US).  <b>(74) Agents:</b> BLOSSER, G., Harley et al.; Senniger, Powers, Leavitt & Roedel, 16th floor, One Metropolitan Square, St. Louis, MO 63102 (US).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> CYTOKININ OXIDASE  <b>(57) Abstract</b>  An isolated protein which exhibits cytokinin oxydizing activity selected from the group consisting of SEQ. ID No. 1, a protein having an amino acid sequence which includes the amino acid sequence of SEQ. ID No. 1, a protein having an amino acid sequence which includes a portion of the amino acid sequence of SEQ. ID No. 1, the included portion being at least about 20 amino acid residues in length and conferring the cytokinin oxidizing activity on the protein, and a protein including an amino acid sequence with at least about 65 % sequence identity to SEQ. ID No. 1, the remainder of amino acid residues being conservatively substituted. Nucleic acids encoding proteins which exhibit cytokinin oxidizing activity and related products and methods are also disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## CYTOKININ OXIDASE

BACKGROUND OF THE INVENTION

The present invention relates to a purified plant cytokinin oxidizing enzyme (*ckx1*) from *Zea mays*, the complete amino acid sequence of which has been elucidated, and to isolated nucleotide sequences encoding the enzyme. The invention further relates to novel methods for moderating the concentration of the enzyme and similar enzymes in plants in order to affect plant cell growth and death. Applications of the invention include the regulation of the production of *ckx1* in plant roots to affect pathogenesis, the regulation to alter plant habit, and the bulk production of *ckx1* enzyme for use in a plant biochemical assay.

Plant cytokinins are a class of plant hormones which, when combined with auxin, control cell division, promote shoot development from callus, release lateral buds from dormancy, and regulate plant structure and growth in a variety of ways. The naturally occurring active cytokinins in most higher plants are free-base zeatin (6-(4-hydroxy-3-methylbut-trans-2-enylamino)purine) (hereinafter Z), and its 9-riboside (hereinafter ZR). Plant tissues normally contain, therefore, Z, ZR, and smaller amounts of N<sup>6</sup>-( $\Delta^2$ -isopentenyl)adenine (hereinafter, iP) derived from biosynthetic precursors. Elevated cytokinin levels are associated with the development of seeds in higher plants, and have been demonstrated to coincide with maximal mitotic activity in the endosperm of developing maize kernels and other cereal grains. Exogenous cytokinin application (via stem injection) has been shown to directly correlate with increased kernel yield in maize. In addition, plant cells transformed with the *ipt* gene from *Agrobacterium tumefaciens* (encoding a

dimethylallylpyrophosphate:5'-AMP transferase capable of increasing cellular production of Z and ZR) showed increased growth corresponding to an increase in endogenous cytokinin levels upon induction of the enzyme. Thus, given the biosignificance of cytokinins to the growth of plants, the ability to manipulate cytokinin levels in higher plant cells is of great commercial and scientific interest.

The action of cytokinin oxidase is a major method of effective cytokinin catabolism in plant cells. This inactivation of cytokinin is accomplished by the oxidative removal of the side chain from cytokinin free bases (or their ribosides) in the presence of molecular oxygen. An example of this reaction with iP is shown in Figure 1a. Although the exact chemical mechanism for this reaction is unknown, it is suspected that the enzyme is reduced during the deprotonation of iP to N<sup>6</sup>-( $\Delta^2$ -isopentenylimino)purine. The purine is then hydrolyzed into adenine and intermediate 3-methyl-2-butenal (Figure 1b).

While the electron acceptor responsible for reoxidizing the reduced enzyme in plant cells is not known, molecular oxygen can do so *in vitro*. Alternatively, the reduced enzyme may be reoxidized *in vitro* by intermediates such as Cu<sup>+2</sup>/imidazole complexes or the artificial electron acceptor dichlorophenolindophenol (DCPIP).

Cytokinin oxidases are known to remove cytokinins from plant cells after cell division, and have also been postulated to be involved in the senescence process. Cytokinin oxidase activities have been shown to positively correlate to the mitosis of endosperm cells in maize kernels, along with the increase in natural cytokinin concentrations. Oxidase activity increases shortly after the increase in endogenous cytokinin levels. A similar correlation was demonstrated with artificially increased cytokinin levels in transgenic

tobacco. Thus, expression of cytokinin oxidases is thought to be involved in the maintenance of hormonal homeostasis in developing plant cells. Because cytokinin oxidases appear to be substrate-inducible, they act in a negative regulatory fashion to reduce elevated cytokinin levels back to basal values. This substrate induction of cytokinin oxidase activity is a significant barrier to potential commercial applications which attempt to manipulate cytokinin levels in transgenic plants through increased cytokinin production.

Cytokinin oxidases have been discussed for a number of plant species, including *Vinca rosea*, beans (*Phaseolus vulgaris* and *lunatus*), wheat (*Triticum aestivum*), tobacco (*Nicotiana tabacum*), *Dianthus caryophyllus*, soy (*Glycine max*), and maize (*Zea mays*). All of these plant cytokinin oxidases have a similar substrate preference for iP and Z, but show limited or no reactivity with bulky, reduced, or aromatic side chain cytokinins. All also exhibit enhanced activity in the presence of copper plus imidazole. However, these enzymes show substantial variation in both specific activity and molecular weight. This is thought to be linked to the occurrence of glycosylated and unglycosylated variants of the protein, both between and within species.

In the case of the glycosylated cytokinin oxidase, the heavily glycosylated protein may present a carbohydrate-rich surface, preventing antibody formation against peptide epitopes. The glyco-epitopes to which antibodies are raised under these conditions are non-specific, and may prevent isolation of the protein, or clones containing the gene which encodes it, via immuno-chromatography or other immunology-based means. An earlier reported attempt to isolate the gene for maize

cytokinin oxidase (*ckx1*) by immunoscreening of maize cDNA library expression products (Burch, 1992) was unsuccessful.

5 As demonstrated, the full amino acid sequence and encoding DNA for a cytokinin oxidase has been a long sought after goal in modern plant physiology.

#### SUMMARY OF THE INVENTION

10 It is therefore an object of the present invention to provide a means by which recombinant cytokinin oxidase may be produced in quantity so that the effects of cytokinin oxidase on plant growth and metabolism may be studied. It is also an object of the present invention to provide a means for the modification of cytokinin oxidase production in plant cells, *in vivo*, in order to modulate  
15 the endogenous cytokinin level of plant cells to effect altered pathogen resistance and plant growth properties.

The present invention, therefore, is directed to a novel, isolated and substantially purified plant cytokinin oxidizing enzyme, (*ckx1*), having a molecular  
20 weight most preferably of about 60 kD, a sequence length of from about 505 to 565 amino acid residues, preferably 525 to 545 amino acid residues, and most preferably 534 amino acid residues, and having cytokinin inactivating activity. The present invention is also directed to a  
25 protein having an amino acid sequence which includes the amino acid sequence of *ckx1* (SEQ. ID NO. 1). The invention is directed as well to a protein which has cytokinin inactivating activity and which includes a portion of the amino acid sequence of *ckx1* at least about  
30 20 amino acid residues in length, where the included portion of the *ckx1* sequence confers the cytokinin inactivating activity on the protein. The invention is directed to proteins which have cytokinin inactivating activity and have at least about 65% sequence identity to

*ckx1* and most preferably at least about 95% sequence identity to *ckx1*, with the remaining amino acids being conservatively substituted.

The invention is directed, moreover, to  
5 substantially isolated nucleic acid polymers encoding *ckx1* or a cytokinin oxidizing homolog thereof. The nucleic acid polymer most preferably has a nucleic acid sequence of SEQ. ID NO. 3 or the predictable variants thereof described in SEQ. ID NO. 10. The invention is  
10 also directed to a substantially isolated nucleic acid polymer which contains a portion of SEQ. ID NO. 2, SEQ. ID NO. 3, or a nucleic acid polymer described by SEQ. ID NO. 10, the portion being at least 60 bp in length. In addition, the invention is directed to nucleic acid  
15 polymers which are able to hybridize with SEQ. ID NO. 2, SEQ. ID NO. 3, or a nucleic acid polymer described by SEQ. ID NO. 10, under conditions of 0.5X to 2X SSC buffer, 0.1% SDS, and a temperature of 55-65°C. Nucleic acid polymers which encode cytokinin oxidases and meet the  
20 above requirements encode proteins of sufficient similarity to *ckx1* to be generally recognized as equivalents of *ckx1* among those skilled in the biochemical arts.

The invention is also directed to a host cell  
25 incorporating a vector containing the aforementioned DNA, and to a method for producing *ckx1* or a homolog thereof using such a host cell. The method preferably comprises first ligating DNA encoding the aforementioned *ckx1* or a segment or homolog thereof and an appropriate promoter  
30 (such as the *RB7* root-specific promoter, Conkling, M.A., et al., U.S. Pat. No. 5,459,252; or *CaMV35S* promoter, Odell, 1985), or a combination of promoters (Hoffman, Patent No. 5,106,739) 3) into an appropriate DNA vector (for instance, pBIN19 for use in *Agrobacterium*  
35 *tumefaciens*). The vector construct may then be directly

transformed into a host cell, such as *Pichia pastoris* (described in Example 2). It may also be incorporated into a secondary vector for transformation into a host cell, such as *Agrobacterium tumefaciens*, and transformed  
5 into a plant cell host (described in Example 4 with *Nicotiana tabacum*).

Alternatively, for production of larger amounts of the enzyme, the DNA encoding *ckx1*, or a portion thereof, may be transformed into *Pichia*, according to the methods  
10 described in Example 2, or in Su, et al., 1996 and Skory, et al., 1996. When transformed into *Pichia spp.*, *ckx1* is secreted into the culture medium because of the presence of a secretory signal peptide at the N-terminus of the *ckx1* coding region. Thus, active enzyme may be readily  
15 purified from bulk *Pichia* cultures without a lysing step. *ckx1* produced in such a manner may be used in a biochemical assay to determine unknown concentrations of cytokinin in biological samples, according to the method of Example 3.

20 A plant host cell generated by the above method may be regenerated into an entire plant. Depending on the promoter used in the vector construct, *ckx1* may be produced constitutively or by induction through natural or artificial environment factors. Plants transformed  
25 with vectors containing tissue-specific or trauma-specific promoters and a sequence encoding *ckx1* can exhibit altered resistance to certain cytokinin-linked plant pathologies, such as infection by certain nematodal or fungal species.

30 The discoveries described herein provide an important analytical tool for, and a critical link in, the development of methods by which the manipulation of cytokinin oxidase activity may be used to either inhibit or enhance a variety of cell growth functions in plants  
35 in a desired manner. Possible uses include the

development of commercial plants with increased grain production, disease resistance, or with more desirable secondary growth characteristics. The enzyme and its encoding nucleic acids have important uses in the study of plant cell growth cycles and senescence. In addition, many other pharmaceutical and agricultural uses for *ckx1* and its gene may be discovered. The enzyme, methods of expressing the enzyme, and methods for its use are described in greater detail below.

Other features and objects of the present invention will be in part apparent to those skilled in the art and in part pointed out hereinafter.

BRIEF DESCRIPTION OF THE FIGURES,  
SEQUENCE IDENTIFICATIONS, AND DEFINITIONS

The invention is further disclosed and illustrated by the accompanying figures.

FIG. 1a & b show an exemplary reaction catalyzed by cytokinin oxidase (Brownlee, 1975), and its putative mechanism as described above, based on Hare, 1994.

FIG. 2 shows agarose gel electrophoresis of RT-PCR DNA fragments which demonstrate that the introns of the *ckx1* gene have been correctly identified.

FIG. 3 shows the standard spectrophotometric absorbance curve (590 nm) obtained when using *ckx1* to assay cytokinin concentrations in solution.

FIG. 4 is a diagram of DNA plasmid pJL7.

FIG. 5 is a diagram of DNA plasmid pBI121.

FIG. 6 is a diagram of DNA plasmid pROM8.

FIG. 7 is a diagram of DNA plasmid pROM9.

FIG. 8 is a diagram of DNA plasmid pROM22.

FIG. 9 is a diagram of DNA plasmid pROM24.

FIG. 10 is a diagram of DNA plasmid pROM26.

FIG. 11 is a diagram of DNA plasmid pROM28.

FIG. 12 is a diagram of DNA plasmid pROM29.

FIG. 13 is a diagram of DNA plasmid pROM30.

FIG. 14 is a diagram of DNA plasmid pROM32.

FIG. 15 is a diagram of DNA plasmid pROM43.

All Sequence Identification abbreviations of amino acids and nucleotides conform to USPTO and WIPO standards.

5 SEQ. ID NO. 1 lists the amino acid sequence of naturally occurring *ckx1* derived from *Zea mays*. This sequence was predicted from the genomic DNA derived nucleotide sequence encoding *ckx1*.

10 SEQ. ID NO. 2 lists the genomic DNA sequence encoding *ckx1*, including introns.

SEQ. ID NO. 3 lists the coding DNA sequence for *ckx1*, with introns excluded. This sequence has been reconstructed in pROM22 of Example 2.

15 SEQ. ID NO. 4 lists the amino acid sequence of an internal tryptic digest fragment of *ckx1*.

SEQ. ID NO. 5 lists the amino acid sequence of an internal tryptic digest fragment of *ckx1*.

20 SEQ. ID NO. 6 lists the nucleic acid sequence of a degenerate DNA probe used to isolate the genomic DNA encoding *ckx1*, as described in Example 1. Note that the residues designated "n" in the sequence are the artificial base inosine (I). Normal conventions have been followed for the indication of degeneracies in the sequence.

25 SEQ. ID NO. 7 lists the nucleic acid sequence of a degenerate DNA probe used to isolate the genomic DNA encoding *ckx1*, as described in Example 1. Note that the residues designated "n" in the sequence are the artificial base inosine (I). Normal conventions have been followed for the indication of degeneracies in the sequence.

30 SEQ. ID NO. 8 lists the nucleic acid sequence of a degenerate DNA probe used to isolate the genomic DNA

encoding *ckx1*, as described in Example 1. Note that the residues designated "n" in the sequence are the

artificial base inosine (I). Normal conventions have been followed for the indication of degeneracies in the sequence.

5 SEQ. ID NO. 9 lists the nucleic acid sequence of a degenerate DNA probe used to isolate the genomic DNA encoding *ckx1*, as described in Example 1. Note that the residues designated "n" in the sequence are the artificial base inosine (I). Normal conventions have been followed for the indication of degeneracies in the  
10 sequence.

SEQ. ID NO. 10 lists the degenerate DNA sequence encoding *ckx1*. As is well known in the art, the several DNA molecules indicated by this group encode a protein with the amino acid sequence of SEQ. ID NO. 1, also known  
15 as *ckx1*. This group follows the conventional rules of degeneracy in the genetic code. Special modifications necessary for expression in certain organisms which do not follow these conventions could easily be made by an individual of ordinary skill in the art.

20 SEQ. ID NO. 11 lists the sequence of a synthetic primer used in the PCR removal of introns in example 2.

SEQ. ID NO. 12 lists the sequence of a synthetic primer used in the PCR removal of introns in example 2.

25 SEQ. ID NO. 13 lists the sequence of a synthetic primer used in the PCR removal of introns in example 2.

SEQ. ID NO. 14 lists the sequence of a synthetic primer used in the PCR removal of introns in example 2.

SEQ. ID NO. 15 lists the sequence of a synthetic primer used in the PCR removal of introns in example 2.

30 SEQ. ID NO. 16 lists the sequence of a synthetic primer used in the PCR removal of introns in example 2.

SEQ. ID NO. 17 lists the sequence of a synthetic linker construct used in example 2.

35 SEQ. ID NO. 18 lists the sequence of a synthetic linker construct used in example 2.

SEQ. ID NO. 19 lists the sequence of a synthetic primer used in PCR to obtain the tobacco RB7 promoter in example 4.

5 SEQ. ID NO. 20 lists the sequence of a synthetic primer used in PCR to obtain the tobacco RB7 promoter in example 4.

10 As used herein, a "substantially purified protein" means that the protein is separated from a majority of host cell proteins normally associated with it or that the protein is synthesized in substantially purified form, such synthesis including expression of the protein in a host cell from a nucleic acid polymer exogenously introduced into the cell by any suitable gene-therapy delivery means.

15 A "substantially isolated nucleic acid polymer" means that the mixture which comprises the nucleic acid polymer of interest is essentially free of a majority of other nucleic acid polymers normally associated with it. A "nucleic acid polymer" includes a polymer of  
20 nucleotides or nucleotide derivatives or analogs, including for example deoxyribonucleotides, ribonucleotides, etc. Genomic DNA, cDNA and mRNA are exemplary nucleic acid polymers.

25 The terms "regulate transcription," "modify transcription," "regulate production," and "modify production," is intended to include promotion and/or repression of transcription or mRNA or production/translation of a protein.

30 The term "expression regulatory sequence" means a nucleic acid polymer sequence ligated to a protein encoding sequence which, when introduced into a host cell, either induces or prevents expression of that protein. These sequences may or may not also encode

proteins used in their regulatory mechanism. Examples of expression regulatory sequences include the *CaMV* promoter, the *ocs* terminator, and the *tet* operator sequences.

5       The term "gene" is intended to include both endogenous and heterologous genes, and specifically, both genomic DNA which encodes a target protein in a naturally occurring cell, and also cDNA encoding the target protein, for example, wherein the cDNA is a part of a  
10       nucleic acid construct such as a plasmid vector or virus which has been introduced into a cell or a cDNA produced by RT-PCR.

      The term "vector" is intended to include any physical or biochemical vehicle containing nucleic acid  
15       polymers of interest, by which those nucleic acid polymers are transferred into a host cell, thereby transforming that cell with the introduced nucleic acid polymers. Examples of vectors include DNA plasmids, viruses, particle gun pellets, and bacteria such as  
20       *Agrobacterium tumefaciens*. The term "primary vector" is intended to mean the first vector used in a transformation series, either as one step (e.g. a plasmid used to transform a yeast cell), or with a "secondary vector" (e.g. a plasmid used to transform *Agrobacterium*  
25       *tumefaciens*, which is later used to transform a plant cell).

      The term "host cell" is intended to mean the target cell for vector transformation, in which the transferred nucleic acid polymer will be replicated and/or  
30       expressed.

      The term "conservative substitution," in the context of amino acid sequences, means the substitution of one amino acid in the sequence with another with a side chain of similar size and charge. An example of a conservative  
35       substitution would be substituting glutamine for asparagine. Conservative substitutions in a protein

sequence which would be expected to have minimal to no impact on protein structure or function can be readily devised by a person of ordinary skill in the biochemical arts.

5       The term "plant products" means any cellular material produced by a plant, especially those which may be used for propagation of the plant. Plant products include seeds, rhizomes, leaves, meristem, roots, and buds.

10       The term "SSC buffer" means a solution of 8.765 g NaCl and 4.41 g sodium citrate in 1 liter of water, pH adjusted to 7.0 by titrimetric addition of 10 N NaOH solution.

15       The contents of each of the references cited herein are being incorporated by reference in their entirety.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to newly sequenced cytokinin oxidase isolated from the maize plant, *Zea mays*, designated *ckx1*, which exhibits a substrate  
20       specific cytokinin oxidizing activity. The enzyme has been linked to the development and maturation of kernels in maize, as well as the senescence of plant tissues. The control of these plant functions appears to be achieved by the degradation of endogenous and exogenous cytokinin  
25       concentrations. Because *ckx1* efficiently oxidizes unsaturated-side-chain free cytokinins and their ribosides, such as iP, Z, and ZR (which also induce production of the enzyme when applied exogenously or endogenously), it is thought that *ckx1* rigorously  
30       controls the availability of these active cytokinins as a part of plant growth regulation.

During the course of developing the claimed invention, the applicant encountered several difficulties which may explain why the search for a cytokinin oxidase  
35       gene has been heretofore unsuccessful. The protein is

expressed at very low levels, and only at certain periods of the plant growth cycle. Therefore, screening a  $\lambda$ -cDNA maize library in E. coli may not yield *ckx1* simply because the gene was not expressed, or not expressed in detectable amounts, when the library was taken. Also, isolation of the protein may have been a problem for some researchers, since its active form does not survive gel filtration. Only through the development of his own screening process, described in Example 1, was the applicant able to isolate the protein in sufficient purity for tryptic digestion and subsequent sequencing.

Even after the protein was purified, obtaining the gene sequence for *ckx1* proved to be difficult.

Apparently, the N-terminal end of *ckx1* is blocked, so a straightforward Edman degradation determination of its N-terminal sequence was impossible. Only small, internal sequences were determinable after tryptic digestion of the protein, SEQ. ID NO. 4 & 5. Because the location of these fragments in the protein's amino acid sequence was unknown, several problems had to be overcome in order to successfully probe the *Zea mays* genome for the *ckx1* gene, as detailed in Example 1. The small size of the sequenced peptides necessitated using a "bookended" criteria (one probe at either end of the replicated DNA) in order to eliminate non-*ckx1* DNA from either side of the *ckx1* gene. One could be reasonably certain that the DNA between two probes would be part of the *ckx1* gene. A hybridization/amplification product size criteria of 300 bp was also necessary in order to distinguish between dimerized degenerate primers and PCR products. A simple degenerate strategy was unlikely to work because of the high degeneracy inherent in the particular amino acid sequences available from the tryptic peptides. Initially, when a standard degenerate nucleotide strategy was used, utilizing all possible oligonucleotides encoding the interior amino acid sequences, the low concentration of

specifically binding primer was not sufficient for the initiation of PCR amplification. The applicant was able to overcome this problem only by using inosine containing degenerate probes with a broader specificity. Also,  
5 several of the degenerate probes were too close together to yield products of the target size, or not in the right order in the gene to produce "bookended" products. Only probes SEQ ID NO. 6, 7, 8, & 9 proved to be of use in isolating the *ckx1* gene.

10 The identity of the isolated gene was verified by two independent methods. First, it was verified by testing the affinity of goat antisera to an *Escherichia coli* produced translation of SEQ. ID NO. 3 for *ckx1* in maize kernel extracts, as per Example 1. Second, it was  
15 confirmed by expressing SEQ. ID No. 3 in *Pichia pastoris* resulting in the secretion of active cytokinin oxidase as per Example 2.

SEQ. ID NO. 2 lists the complete genomic DNA sequence of *ckx1*, which yields the glycosylated form of  
20 the protein when expressed in *Zea mays*. The predicted location of the introns in the genomic sequence was verified using the reverse transcriptase- polymerase chain reaction to find the actual transcribed length of RNA, as per Example 1. SEQ. ID NO. 3 lists the coding DNA  
25 sequence for *ckx1*, which yields the amino acid sequence set forth in SEQ. ID NO. 1.

The state of the art of molecular biology is now sufficiently advanced that minor alterations can be made to a DNA sequence with relative ease and precision. A  
30 moderately skilled laboratory technician can follow the directions of one of the commercially available site-directed mutagenesis kits (for instance, the GeneEditor™ offered by Promega Corp., Madison, Wisconsin) to effect any number of changes to a DNA nucleotide sequence. Also  
35 well known are the general rules governing the genetic code, by which triplet nucleotide codons are translated

into an amino acid sequence by standard biochemical processes. Thus, the applicant considers the group of DNA sequences denoted by the consensus sequence of SEQ. ID NO. 10, which code for the amino acid sequence of *ckx1* in SEQ. ID. NO. 1, to be within the present invention. Although some variation in the genetic code and the GC% content occurs amongst some phyla, the rules governing these variations have also been well documented, and are within the reasonable skill of one versed in the molecular genetic arts. Thus, the applicant also considers any other nucleic acid sequence which encodes the amino acid sequence SEQ. ID NO. 1 to be within the scope of the present invention.

In addition, although the understanding of the field of protein biochemistry is not as complete as that of molecular genetics, the person of ordinary skill in the art of biochemistry is capable of predicting, with reasonable certainty, when certain substitutions to the primary amino acid sequence structure of a protein will not result in any appreciable modification of a protein's structure or function. Such conservative substitutions are made by replacing an amino acid in the sequence with another containing a side chain with like charge, size, and other characteristics. For instance, the amino acid alanine, which has a small nonpolar methyl side chain, generally can be replaced by glycine, an amino acid which has a small nonpolar hydrogen side chain, without any noticeable effects. Likewise, the amino acid asparagine, with a moderately bulky, polar ethamide side chain, usually can be replaced with glutamine, which has a moderately bulky, polar propamide side chain, without noticeable effects. To the extent that such conservative substitutions can be made while retaining 65%, preferably 80% or more identity to SEQ. ID NO. 1 and cytokinin oxidizing activity, such altered proteins are within the scope of the present invention.

Cytokinin oxidases are known to exist in a variety of non-glycosylated and glycosylated forms in several species of higher plants, including maize. This modification is thought to be involved in compartmentalizing cytokinin oxidases for various uses inside and outside the cell. The extent of glycosylation of the enzyme may also account for the wide variety of molecular weights observed between the cytokinin oxidases of various species. However, because substrate specificity, a requirement of molecular oxygen for activity, and copper concentration reaction rate effects in vitro are highly conserved among all higher plant cytokinin oxidases, a common domain and active site structure is believed responsible for the cytokinin oxidizing activity of all enzymes. Thus, the present invention is also directed to a protein which exhibits cytokinin oxidizing activity and which contains an amino acid sequence at least about 20 amino acids in length which is 90% identical (or would be identical with conservative amino acid substitutions) to a similarly sized portion of SEQ. ID NO. 1.

Another object of the current invention is the regulated production of *ckx1* in various host cells, either for later bulk isolation, or regulated intracellular production. For instance, *ckx1* may be produced in unglycosylated form in prokaryotes, such as *E. coli*, as illustrated in example 1. More preferably, the protein may be produced in eukaryotes, as illustrated by the *Pichia* production of example 2. An added benefit of producing the protein in *Pichia* is that the protein is secreted into the culture media where it may be readily purified. Alternatively, the protein may be produced in higher eukaryotes, more preferably plants, such as the *Nicotiana* constructs of example 4 either as the glycosylated or non-glycosylated forms. Other examples of suitable host plant cells include *Zea mays*,

*Arabidopsis thaliana*, *Brassica spp.*, and *Oryza sativa*. As illustrated, *ckx1* can be produced in plants either in an unregulated fashion, as shown here under the *CaMV* promoter, regulated by an artificial stimuli, as shown here under a tet operator combined with a *CaMV* promoter, and regulated by an environmental stimuli, as shown here under the *RB7* promoter, which induces root-specific production of a protein in response to nematodal attack.

The several aspects of the present invention, including the *ckx1* protein and the nucleic acid polymers which encode it, the cytokinin oxidizing activity of enzymes such as the *ckx1* enzyme, and the regulation of cytokinin levels in plant cells by *ckx1*, collectively enable several practical applications, including both agricultural and research uses.

The applicant has devised an application for bulk cytokinin oxidase which greatly facilitates plant physiology research. Cytokinin oxidase can be reoxidized after oxidizing cytokinins by the synthetic oxidizing agent dichlorophenolindophenol (DCPIP). DCPIP demonstrates a reduced absorbance at 590 nm upon reduction, which can be spectrophotometrically quantified. The cytokinin concentration in a sample may be determined indirectly by measuring the decrease in oxidized DCPIP as *ckx1* oxidizes the cytokinins present in the sample. More sensitive redox dyes are also available to increase the sensitivity of the assay. Therefore, it is another object of the present invention to provide a simple, fast, and effective means to assay cytokinin concentrations in a sample.

The applicant has also devised an application for the modified production of *ckx1* in plants. Cytokinins are associated with several types of plant pathogenesis, including the formation of nematode feeder cells, and.

5 fungal invasion of plant tissues. Thus, the pathogen-exposure-regulated production of *ckx1* can modify the efficacy of pathogenesis through cytokinin-utilizing mechanisms such as that utilized by the root-knot nematode, *Meloidogyne* spp. (Bird, et al., 1980), or fungal species such as *Ustilago maydis*. Thus, it is an object of the present invention to provide a method of moderating cytokinin-mediated pathogenesis by transforming a plant with a *ckx1* producing DNA construct.

10 The following examples illustrate the principles and advantages of the invention.

#### EXAMPLES

15 Example 1: Isolation of *ckx1* and characterization of its encoding sequence

Maize kernels were chosen as the raw material for purification because of the relatively high concentrations of cytokinin oxidase present about a week after pollination. Using the procedure set forth below, approximately 1.66  $\mu$ g protein per kg maize yield may be obtained. Field grown maize (Pioneer 3180 or 3379) was hand pollinated, and immature ears were harvested between 5 and 8 days later (Dietrich, 1995). Kernels were harvested immediately, and frozen at  $-80^{\circ}\text{C}$  until extraction. After being powdered in liquid nitrogen, the kernels were blended in 1 kg lots with 1700 ml of Buffer A (50 mM Tris, 5 mM EDTA, ascorbic acid 0.4% w/v; 10 mM  $\beta$ -mercaptoethanol, pH 8.5). Acid washed PVPP (200 g wet weight, equilibrated w/ Buffer A) was stirred in immediately. The slurry was filtered through Miracloth and centrifuged at 23,500 x g for 15 minutes to remove debris. Polyethyleneimine solution (5% v/v, pH 8.5) was added dropwise to the centrifuged supernatant, to a final concentration of 0.05%. After recentrifugation at 23,500

x g for 10 minutes, the supernatant was filtered through a 600 g pad of PVPP (prepared as above). Ammonium sulfate fractionation was performed, and the protein precipitating between 40% and 65% saturation was collected by centrifugation and dissolved in a minimum volume of Buffer B (10 mM Tris, 1 mM EDTA, 1 mM  $\beta$ -mercaptoethanol, pH 8.5). Insoluble material was removed by centrifugation at 35,000 x g for 20 minutes. At this point, glycerol may be added to the supernatant to 10% v/v, allowing the protein to be stored at -80°C indefinitely without loss of activity.

After dialyzing the supernatant from the ammonium sulfate fractionation against Buffer B, the fraction was applied to a DEAE-cellulose column (Whatman DE-52, 500 ml, available from the Whatman Group, Clifton, New Jersey) at the rate of 10 ml/minute. After washing w/ buffer B, the column was eluted with a linearly increasing concentration of KCl in Buffer B, to 200 mM over 600 ml. Protein content was measured using the Bradford dye binding assay (Bradford, 1976). Fractions were analyzed for oxidase activity as described below.

Two assays were used to screen purification steps for cytokinin oxidase activity. The first was the Schiff base formation assay measuring the production of dimethylallylaldehyde from iP, as described by Libeross-Minotta (1995), which was used up through the DEAE column step. A second assay, developed by the applicant, was used in the remaining purification steps. In this assay, the transfer of reducing equivalents from isopentenyladenine to dichlorophenolindophenol (DCPIP), catalyzed by *ckx1*, allows the observance of reactivity by measuring absorbance change at 590 nm. In a final volume of 250  $\mu$ l, the assay contains: 100 mM Phosphate buffer, pH 7.0; 1.0 mM EDTA; 0.05 mM DCPIP; 0.1 mM iP; 100  $\mu$ g/ml BSA; and the sample tested. After the addition of the enzyme, absorbance change is read at 590 nm for 10 minutes.

After purification on the DEAE column, the major pooled active material was dialyzed against Buffer C (20 mM Tris, 0.5 M NaCl, 1.0 mM CaCl<sub>2</sub>, pH 7.4), and applied to a concanavalin A agarose column (100 ml, from Sigma Aldrich, St. Louis, Missouri) and washed w/ Buffer C (270 ml). Glycosylated proteins were eluted with a step gradient of buffer C containing  $\alpha$ -D-methylmannoside to 1 M over 400 ml. The relatively long retention time when eluted under these conditions indicates that the glycosylated form of *ckx1* was isolated. Active fractions from the lectin-affinity chromatography were then dialyzed against Buffer D (10 mM Tris, 1 mM EDTA, pH 8.5) and applied to a high resolution anion exchange column (FPLC MonoQ, 1 ml, from Amersham Pharmacia Biotech, Ltd., San Francisco, California), and eluted with a linear gradient of KCl to 0.12 M over 24 minutes at 1 ml/minute. The active fractions from the ion exchange column were then concentrated, dialyzed against Buffer B, brought to 1.5 M ammonium sulphate and applied to a hydrophobic interaction column (FPLC phenylsuperose, 1 ml, Pharmacia) equilibrated in Buffer B containing 0.6 ammonium sulfate. After washing with 0.6 M ammonium sulfate for 25 minutes, the concentration of ammonium sulfate was reduced successively to 0.45 M over 15 minutes and then to zero over 60 minutes.

Native gel electrophoresis was performed as illustrated in Ornstein (1964) and Davis (1964). Gels were then stained for cytokinin oxidase activity by the DCPIP procedure described above. Enzyme activity was revealed as a transparent band against a blue background. Denaturing SDS polyacrylamide gel electrophoresis was then carried out as illustrated in Laemmli (1970). When testing fractions for homogeneity, the gel was stained as described by Møller (1995). At the final purification step, the enzyme was stained with Coomassie Blue R250. The purified protein was analyzed by tryptic digestion, HPLC separation of digest, and Edman degradation

sequencing of the tryptic polypeptides. Several polypeptide sequences were obtained from this analysis, including SEQ. ID NOS. 4 and 5. From these sequences, reverse translation primer probes, SEQ. ID NOS. 6, 7, 8, and 9, were devised, with inosine substituted at highly degenerative positions. Primers SEQ. ID NOS. 6 and 9 were then combined with maize genomic DNA, and hot-start touchdown PCR was performed (Ault, 1994) for 40 cycles. PCR products were separated on agarose gel, and an approximately 440 bp fragment was chosen to use as a hybridization probe. The identity of the fragment was confirmed by showing that it could be amplified by PCR with the nested internal primers SEQ. ID NOS. 7 and 8. The fragment amplified by the primers SEQ. ID NOS. 6 and 9 was then ligated into linearized pCRII DNA and transformed into *E. coli* (INVαF', Invitrogen Corp., Carlsbad, California). After cloning and reisolation of DNA, plasmid inserts were sequenced using the Prism dideoxy terminator method of Applied Biosystems, Foster City, CA, to verify that the sequenced tryptic digest polypeptides were encoded by the fragment.

Once the large fragment had been verified, it was labeled with  $^{32}\text{P}$  by primer extension using the Klenow fragment of DNA polymerase and primers SEQ. ID NOS. 6 and 9. Maize genomic library phage (in λ-FIXII, from Stratagene, La Jolla, California) were diluted in SM Buffer, and appropriate numbers added to freshly prepared *E. coli* (XL1-Blue MRA (P2), Stratagene) in 10 mM  $\text{MgSO}_4$  and incubated at 37°C for 15 minutes. NZY top agar at 48°C was added and the mixture plated onto NZY agar plates. After incubation for approximately 8 hours at 37°C, plates were cooled to 4°C for 2 hours and phage were adsorbed onto sheets of Hybond N membrane (Amersham Pharmacia Biotech, San Francisco, California). Membranes were air-dried for 10 minutes, and incubated successively with 0.5 M NaOH plus 1.5 M NaCl, 500 ml, 5 minutes; 0.5 Tris-Cl pH 8.0 plus 1.5 M NaCl, 500 ml, 5 minutes; and 2 x SSC, 500 ml,

5 minutes. They were blotted dry and baked at 80°C until dry (approximately 15 minutes). Membranes were pre-hybridized at 45°C for less than one hour in 50% formamide, 5 x SSPE, 2 x Denhardt's solution, 0.2% SDS, 5 100-200 µg/ml denatured herring sperm or calf thymus DNA (Maniatis, 1990).

The labeled DNA fragment was then denatured at 100°C for 5 minutes, added to the pre-hybridization solution, and hybridized for 16 hours at 45°C. For the primary 10 screen, phage were plated at a density of 500 pfu/cm<sup>2</sup> and membranes were washed at high stringency. For subsequent plaque purification, candidate phage were plated at a density of 70 pfu/cm<sup>2</sup> and 2 pfu/cm<sup>2</sup> and membranes were washed at medium stringency. After three rounds of 15 purification, subfragments of the insert were removed from positive phage DNA by restriction and subcloned into pBluescript (Stratagene) for characterization by restriction digestion and sequence analysis. Two overlapping plasmid subclones, pROM2 (a HindIII insert) 20 and pROM3 (an Xho-BamHI insert) each contained part of the gene for *ckx1*. These overlapping sections have been fused into the plasmid clone pROM10. The plasmids pROM2, pROM3, and pROM10 have been deposited with the American Type Culture Collection as ATCC Nos. 209573, 209572, and 25 209571, respectively. The sequence of the cloned DNA in pROM2 and pROM3 provided the genomic sequence of *ckx1*, SEQ. ID NO. 2, which was verified by the inclusion of sequences coding for the tryptic digest fragments obtained above.

30 The location of the introns in the *ckx1* gene, and the coding sequence for *ckx1* (SEQ. ID NO. 3) was verified by use of the reverse transcriptase polymerase chain reaction (RT-PCR) according to the following procedure.

35 Total RNA (2-4 µg, DNAase I-treated) from kernels harvested five days after pollination (5 DAP) was primed for RT-PCR using oligonucleotides bracketing the intron

sites. Primers 2031f (*ckx1* genomic sequence 2031-2050) and XBH1 (reverse complement of *ckx1* genomic sequence 2553-2570) covered the first intron site. Primers 3160f (*ckx1* genomic sequence 3160-3179) and 3484r (reverse complement of *ckx1* genomic sequence 3465-3484) covered the second intron site. Reverse transcription was at 50°C for 50 min. with 200U Superscript II (Gibco-BRL) in 20 µL total volume with 25 mM Tris-Cl, pH 8.3, 37.5 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.5 mM dNTPs, 5mM dithiothreitol, 40U RNAasin (Promega, Madison, Wisconsin). PCR was performed with 4-10% of the RT reaction product as template and 1U Taq polymerase in 1X PCR buffer (Roche Diagnostics, Basel, Switzerland), 200 µM dNTPs, and 0.5 µM each primer. Reaction conditions were: an initial denaturation at 95° C for four minutes followed by thirty-five cycles of 95° denaturation for one minute, 60° annealing for one minute, and 72° extension for one minute. A final four minute extension was carried out at 72°. PCR products were resolved on 1.5% agarose gels. PCR products were excised from gels and sequenced to determine splice site junctions.

RT-PCR of maize kernel RNA (5 days after pollination) with primers designed to span the first and second intron locations demonstrated PCR product sizes consistent with splicing out of these introns in the mature oxidase mRNA. The primers bracketing the first intron should give a 539 bp product if the intron is present and a 127 bp product if it has been spliced out. Likewise, the primers bracketing the second intron should give a 324 bp product if the second intron is present and a 232 bp product if it has been spliced out. As shown in

Figure 2, the PCR products were 127 bp and 232 bp respectively, indicating that the introns had indeed been spliced out. Sequence analysis of the fragments confirmed that splicing had occurred as predicted.

5       The identity of the *ckx1* protein and gene were further verified by the following immunological technique.

Polyclonal antibodies were raised in goats to the peptide produced by expression of a *ckx1* gene fragment in  
10   *E. coli*. This avoided the presentation of a glycosylated surface when raising antibodies to the gene product and the problems encountered by Burch, et al., when raising antibodies to naturally occurring *ckx1*. Goat antibodies from immunized and unimmunized animals were partially  
15   purified by sodium sulfate precipitation (Williams, et al., 1967). Affinity columns were prepared by coupling these purified antibodies (2 mg) to Aminolink Plus gels (1 mL, Pierce) at pH 10, following the manufacturer's protocol. Activity depletion assays were performed by  
20   adding 0.2 mL (27 µg) of a maize ConA-fractionated oxidase preparation plus 0.8 mL phosphate buffered saline (PBS) to each column. The columns were capped and incubated for one hour at room temperature. Eluate was collected and assayed for cytokinin oxidase activity.

25       Antibodies to *ckx1* recognize the major maize cytokinin oxidase enzyme. A column containing immobilized anti-*ckx1*-fragment antibody was able to deplete cytokinin oxidase activity from a ConA-fractionated maize extract. A control column of  
30   unimmunized goat antibodies was not able to do so.

Example 2: Expression of *ckx1* in *Pichia pastoris*

*ckx1* protein may be produced in bulk by the following procedure for use in applications such as the cytokinin assay of Example 3, or application to plant materials.

Expression of *ckx1* in *Pichia pastoris* was carried out in four stages:

Step 1. Removal of the introns from *ckx1*

Step 2. Removal of the maize promoter and construction of the appropriate expression cassette using the intron-less construct

Step 3. Transformation of the final construct into *Pichia* and

Step 4. Expression of *ckx1* in *Pichia*

Removal of the right-most intron was accomplished by splicing by overlap extension (Horton et al., 1989) and for the left-most intron by ligation of suitable restriction fragments. The resulting intronless cassette was inserted into the *Pichia* expression vector pPICZ-A to give the expression construct pROM24 (FIG. 9). This construct was introduced into the requisite *Pichia* host and grown in the presence of methanol to induce *ckx1* expression. Appropriate control *Pichia* lines (containing the vector alone or a recombinant expressing human serum albumin) were grown in parallel.

No cytokinin oxidase activity was observed in *Pichia* cell lysates containing *ckx1* or the controls at any time during the growth curve. However, the *Pichia* line harboring *ckx1* expressed and secreted high levels of cytokinin oxidase activity into the growth medium. No activity was observed in control supernatants. This provided further verification that *ckx1* does indeed encode a cytokinin oxidase.

Step 1. Intron removal: The second intron (SEQ. ID NO. 2 residues 3219-3312) was removed by splicing by overlap extension with selected *ckx1* restriction fragments as templates to limit artifactual priming. The following table lists the primers and templates used.

Intron	Left Primer	Right primer	Template DNA
amplification #1	TGGGAATTCATGGGGAGA TGGTGACGTGCTC (SEQ. ID NO. 11)	GCCGTCCCATGGATTGT TGAGGGGGTAGAC (SEQ. ID NO. 12)	PROM2 <i>Nco</i> I/ <i>Pma</i> I (700 bp) fragment
amplification #2	CTCAACAAATCCATGTGGG ACGACGGCATGTGGCGG (SEQ. ID NO. 13)	GCGGTCTAGATCTAACTAAA ACATGCATGGGCTATCATC (SEQ. ID NO. 14)	PROM2 390 bp <i>Pma</i> I + <i>Vsp</i> I
amplification #3	ATGGGAATTCATGGGGAG ATGGTGACGTGCTC (SEQ. ID NO. 15)	GCGGTCTAGATCTAACTAAA ACATGCATGGGCTATCATC (SEQ. ID NO. 16)	Products from amplifications #1 and # 2

- 10 PCR products from reactions #1 and #2 were gel-purified and used in the final PCR step. The final product from reaction #3 was cloned and sequenced and a *Pfl*MI/*Xba*I subfragment was substituted for the intron-containing *Pfl*MI/*Xba*I subfragment of pROM7. The construct was
- 15 designated pROM19. The first intron (SEQ. ID NO. 2 residues 2113-2524) was then removed from pROM19 by replacement of the DNA between the restriction sites *Pma*I and *Nco*I with a linker constructed from the
- 20 oligonucleotides CCGGTTTTGGTACCGGT (SEQ. ID NO. 17) and CATGACCGGTACCAAAA (SEQ. ID NO. 18). The product was designated pROM20. Extraneous linker-associated bases were removed by digestion with *Pma*I followed by religation. The product, designated pROM22 (FIG. 8), contained the three fused *ckx1* exons and the maize *ckx1*
- 25 promoter.

Step 2. The *Pichia* expression cassette: The maize promoter was removed by partial digestion of pROM22 (FIG. 8) with *Aat*II, filling in the sticky ends with T4 DNA polymerase, and redigestion with *Bgl*II. The exon fusion

(containing *ckx1* and including its putative signal peptide) was ligated into the *Pml1/BsmB1* restriction sites of pPICZ-A (Easy-Select *Pichia* Expression Kit version B, Invitrogen Corp.). The resulting plasmid was designated pROM24 (FIG. 9).

Step 3. Transformation into *Pichia* strain X33 as described in the Invitrogen protocol: The plasmid pROM24 (FIG. 9) (10  $\mu$ g) was digested with *Dra1* and electroporated into competent X33 cells in a 2 mM cuvette at 1.75 kV (GenePulser, Bio-Rad Laboratories, Hercules, California). Selection on YPDS with 100  $\mu$ g/mL zeocin resulted in many colonies. One (PP*ckx1*) was selected for expression studies.

Step 4. Expression of the oxidase: The transformant PP*ckx1* was inoculated into BMGY medium (50 mL) and grown overnight. Cells were pelleted, resuspended in BMMY (containing 0.5% v/v methanol), diluted to an  $A_{600}=1$  and grown at 30°C with vigorous shaking. Additional methanol was added to 0.5% v/v at 24, 48 and 72 hours post-inoculation. Samples were harvested for assay of cytokinin oxidase activity in cell lysates or in culture supernatants. *Pichia* strains X33 (WT, no insert) and GS115, (secreted human serum albumin insert) served as controls.

#### Example 3: Use of Recombinant Cytokinin Oxidase in a Rapid Assay Method for Cytokinin

Cytokinins were measured by mixing 100  $\mu$ L of a buffer mixture containing phosphate buffer (250 mM, pH 7.0), EDTA (2.5 mM) and DCPIP (0.125 mM), and an excess of recombinant cytokinin oxidase with solutions (150  $\mu$ L) of zeatin at various concentrations. The net change in absorbance was measured at 590 nm.

Figure 3 illustrates the change in absorbance when the assay is used to measure the cytokinin zeatin. The method is capable of measuring as little as 2 nmol zeatin but, the major advantage of the assay over the prior art is its rapidity. Assays can be preformed in as little as five minutes, significantly faster than radioimmunoassays (MacDonald and Morris, 1985). Further, the method can be integrated into cytokinin production systems by coupling the *ckx1* gene to such cytokinin producing genes as *ipt* or *tzs*, in order to assay cytokinin production in vitro.

Example 4: Unregulated and Regulated Expression of *ckx1* Constitutively and in the Roots of *Nicotiana tabacum*

The following procedure, which produces tobacco plants altered to express *ckx1* constitutively and in their roots, is a slight modification of the standard protocols described in Draper, et al., 1988.

*Nicotiana tabacum* cultivar Xanthi is a standard tobacco line. Disarmed *Agrobacterium tumefaciens* strains such as LBA4404 (Hoekema, et al., 1985) are used. Murashige and Skoog salts, phytagar, sucrose, etc. are reagent or tissue-culture grade.

Three separate constructs were made with the *ckx1* coding sequence (SEQ. ID NO. 3) to effect three different patterns of *ckx1* expression in transformed tobacco plants. Constructs were based on the BIN19 plasmid primary vector (which includes an *Agrobacterium* compatible replication origin and kanamycin selection markers, Bevan, et al., 1984). The following constructs were made:

*CaMV-ckx1-nos*: The cauliflower mosaic virus promoter with a *nos* enhancer (U.S. Patent No. 5,530,196), are present in pBI121 (FIG. 5) (available from Clontech, Palo

Alto, California). The 8-glucuronidase gene of pBI121 was excised with a *Bam*HI/*Eco*ICrI digest. The coding sequence of *ckx1* (SEQ. ID NO. 3) was obtained by a *Bgl*II digest of pROM26 (FIG. 10) (pROM24 (FIG. 9) altered by site-directed mutagenesis to contain another *Bgl*II and an *Xho*I restriction site). The final construct pROM30 (FIG. 13), which induces strong constitutive expression of *ckx1*, was created by digesting the altered pBI121 with *Bam*HI and ligating in the *ckx1* coding sequence.

10        *CaMV-tet-ckx1-ocs*: The *ckx1* coding region of pROM26 (FIG. 10) was isolated by *Bgl*II digestion and ligated into the *Bam*HI site of pUCA7-TX (Gätz, et al., 1992) to form pROM32 (FIG. 14). The tetracycline-regulated operator element (U.S. Patent No. 5,464,758) from pUCA7-TX and the inserted coding region were excised from  
15        pROM32 (FIG. 14) by *Pvu*II digestion. The final construct pROM29 (FIG. 12), was produced by ligating the excerpt into the *Sal*I site of pJL7 (FIG. 4) (a BIN19 type plasmid). When this construct is introduced into tobacco  
20        previously transformed to express the tet repressor protein, no *ckx1* activity will be expressed until repression is relieved by addition of tetracycline. This construct may be of particular use in host organisms where strong constitutive expression of *ckx1* results in  
25        abnormal growth patterns.

*RB7-ckx1-nos*: The tobacco RB7 root specific promotor (U.S. Patent No. 5,750,386), was obtained by the following method. A fragment containing the promoter was PCR-amplified from tobacco genomic DNA using  
30        GACACCATTCCTCAAGCATACCCC (SEQ. ID NO. 19) and GTTCTCACTAGAAAAATGCCCC (SEQ. ID NO. 20) as primers. This 1400 bp product was ligated into the pCRII plasmid

(Invitrogen) to produce pROM8 (FIG. 6). The *HindIII*-*EcoRI* region of the insert, containing the nematode-specific portion of the promoter, was then excised and ligated into the *HindIII*-*EcoRI* site of pBluescriptII KS+

- 5 (Stratagene) to produce pROM9 (FIG. 7). The *EcoRI*-*NsiI* fragment of pROM24 (FIG. 9), containing the *ckx1* coding sequence, was excised and ligated into the *EcoRI*-*PstI* site of pROM9, to make pROM28 (FIG. 11). The final construct pROM43 (FIG. 15), which induces expression of
- 10 *ckx1* in roots when the transformed plants' roots are attacked by nematodes, was produced by ligating a *HindIII*-*SacI* fragment of pROM28 into the *HindIII*-*SacI* site of pBI121 (FIG. 5).

	Media Component	MSS	PC	SI	MSSK
15	MS Salts (g/l)	4.3	4.3	4.3	4.3
	Sucrose (g/l)	30.0	30.0	30.0	30.0
	Phytagar (g/l)	7.5	5.0	5.0	7.5
	NAA (mg/l)	0	1.0	1.0	0
	BAP (mg/l)	0	0.1	0.1	0
20	Timetin (mg/l)	0	0	200	200
	Kanamycin (mg/l)	0	0	50.0	50.0
	pH	5.7	5.7	5.7	5.7

- 25 *Nicotiana* was continuously maintained by axenic shoot tip culture on MSS and sub-cultured at 4 week intervals.

- 30 The three binary plasmid constructs described above were electroporated into the disarmed *A. tumefaciens* host and transformants were grown under suitable selection (An, et al. 1985). Late-log phase cultures were used for transformation.

Young axenic tobacco leaves (3 to 4 weeks after tip culture) were dissected into 4 to 6 segments, excluding

the largest vasculature. Segments were cultured on PC, abaxial side up, for 48 hours. The leaf pieces were then soaked in transformed *A. tumefaciens* culture diluted with water (1:1), for 1 hour. Pieces were then placed on the original PC medium for 48 hours. The leaves were then washed with water thoroughly and placed on SI medium and removed to fresh SI medium every 7 to 10 days. Although no adverse developmental effects were observed with *Nicotiana* transformants, the use of a non-substrate cytokinin such as benzylaminopurine, or tetracycline repression, may be needed for the culture of some host plants. Shoots were removed onto MSSK media when 1 cm long. Successive tip culture was carried out for 2 to 3 transfers, after which the transgenic plants were maintained on MSS media.

In light of the detailed description of the invention and the examples presented above, it can be appreciated that the several objects of the invention are achieved. The explanations and illustrations presented herein are intended to acquaint others skilled in the art with the invention, its principles, and its practical application. Those skilled in the art may adapt and apply the invention in its numerous forms, as may be best suited to the requirements of a particular use. Accordingly, the specific embodiments of the present invention as set forth are not intended as being exhaustive or limiting of the invention.

BIBLIOGRAPHY

An, G., et al., 1985: "New cloning vehicles for transformation of higher plants," EMBO J. 4:277-84

5 Ault, G., 1994: "Type-specific amplification of viral DNA using touchdown and hot start PCR," J. Virol. Meth. 46:145-56.

Bechtold, N., et al., 1993: "In planta *Agrobacterium* mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants," CR Acad. Sci. Paris  
10 Sciences del la vie/ life sciences 316:1194-99.

Bevan, M., 1984: "Binary *Agrobacterium* vectors for plant transformation," Nuc. Acids Rsrch. 12:8711-21.

Bird, A.F., et al., 1980: "The involvement of cytokinins in a host-parasite relationship between the  
15 tomato (*Lycopersicon esculentum*) and a nematode (*Meloidogyne javanica*)," Parasitology 80:497-505.

Bradford, M., 1976: "A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding,"  
20 Analytical Biochem. 72:248-54.

Brownlee, B.G., et al., 1975: "3-Methyl-2-butenal: An Enzymatic Degradation Product of the Cytokinin, N<sup>6</sup>-( $\Delta^2$ -isopentenyl)adenine," Can. J. Biochem. 53:37-41.

Burch, L.R., et al., 1992: "Cytokinin Oxidase and the Degradative Metabolism of Cytokinins," in Kaminek, et al. Physiology and Biochemistry of Cytokinins in Plants. pp. 229-32, STP Academic Publishing, The Hague, Netherlands.

Chen, H., et al., 1996: "Novel methods of generating specific oligonucleotide inhibitors of viral polymerases." Methods in Enzymology 275:503-20.

Davis, B., 1964: "Disk Electrophoresis II. Method and application to human serum proteins," Ann. N.Y. Acad. Sci. 121:404.

Dietrich, J.T., et al., 1995: "Changes in cytokinins and cytokinin oxidase activity in developing maize kernels and the effects of exogenous cytokinin on kernel development," Plant Physiol. Biochem. 33:327-36.

- 5 Draper, J., et al., 1988: Plant Genetic Transformation and Expression: A Laboratory Manual. Blackwell Scientific Publications.

Gätz, C., et al., 1992: "Stringent repression and homogenous de-repression by tetracycline of a modified  
10 CaMV 35 S promoter in intact tobaccod plants." Plant J. 2(3):397-404.

Hare, P.D., et al., 1994: "Cytokinin oxidase: Biochemical features and physiological significance," Physiologia Plantarum 91:128-35.

- 15 Hoekema, A., et al., 1985: "Non-oncogenic plant vectors for use in Agrobacterium binary systems," Plant. Mol. Biol. 5:85-89.

Horton, R.M., et al., 1989: "Engineering hybrid genes without the use of restriction enzymes: gene  
20 splicing by overlap extension," Gene 77:61-8.

Koziel, et al., 1993: "Field performance of elite transgenic maize plants expressing an insecticidal protein form *Bacillus thuringiensis*," Bio/Technology 11:194

- 25 Laemmli, U.K., 1970: "Cleavage of structural proteins during the assembly of the head of bacteriophage T4," Nature 227:680-685.

Liberos-Minotta, C., et al., 1995: "A colorimetric Assay for Cytokinin Oxidase," Analytical Biochem.  
30 231:339-341.

MacDonald, E.M.S., et al., 1985: "Isolation of cytokinins by immunoaffinity chromatography and isolation by HPLC-radioimmunoassay," Methods in Enzymology 110: 347.

- 35 Maniatis, T., et al., 1990: Molecular Cloning. A Laboratory Manual 2nd Ed. Cold Spring Harbor Laboratory Press.

Møller, H.J., et al., 1995: "Improved method for Silver Staining of Glycoproteins in Thin Sodium Dodecyl Sulfate Polyacrylamide Gels," Analytical Biochem. 226:371-74.

- 5 Odell, et al., 1985: "Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter," Nature 313:810.

- Ooms, G., et al., 1987: "Genetic transformation in two potato cultivars with T-DNA from disarmed  
10 *Agrobacterium*," Theor. Appl. Genet. 73:744-50.

Ornstein, L., 1964: "Disk Electrophoresis I. Background and theory," Ann. N.Y. Acad. Sci. 121:321.

- Potrykus, 1991: "Gene transfer to plants: assessment of published approaches and results," Annu. Rev. Plant  
15 *Physiol., Plant Mol. Biol.*, 42:205.

Singer, B.S., et al., 1996: "Libraries for genomic SELEX," Nucleic Acids Resrch. 25:781-86.

- Skory, C.D., et al., 1996: "Expression and secretion of the *Candida wickerhamii* extracellular beta-glucosidase  
20 gene, bgls, in *Saccharomyces cerevisiae*," Current Genetics 30:417-422.

Smith, C.T.S., et al., 1988: "Antisense RNA inhibition of polygalacturonase gene expression in transgenic tomatoes," Nature 334:724-26.

- 25 Su, W., et al., 1996: "Identification in vitro of a post-translational regulatory site in the hinge 1 region of *Arabidopsis* nitrate reductase," Plant Cell 8:519-27.

- Van der Krol, A.R., et al., 1990: "Inhibition of flower pigmentation by antisense CHS genes: promoter and  
30 minimal sequence requirements for the antisense effect," Plant Molec. Biol. 14:457-66.

Williams, C. A, et al., 1967: Methods in Immunology and Immunohistochemistry I p. 319.

WE CLAIM AS OUR INVENTION:

1. A substantially purified protein which exhibits cytokinin oxidizing activity, selected from the group consisting of:

- (a) SEQ. ID NO. 1,
- 5 (b) a protein having an amino acid sequence which includes the amino acid sequence of SEQ. ID NO. 1,
- (c) a protein having an amino acid sequence which includes a portion of the amino acid sequence of SEQ. ID NO. 1, the included portion being at least about 20 amino
- 10 acid residues in length and conferring the cytokinin oxidizing activity on the protein, and
- (d) a protein including an amino acid sequence with at least about 65% sequence identity to SEQ. ID NO. 1, the remainder of amino acid residues being conservatively substituted.

2. The protein of claim 1 wherein the protein is purified.

3. The protein of claim 1 which is SEQ. ID NO. 1.

4. The protein of claim 1 produced by a host cell transformed with a vector containing a nucleic acid polymer which encodes the protein of claim 1.

5. A substantially isolated nucleic acid polymer encoding a protein of claim 1.

6. The substantially isolated nucleic acid polymer of claim 5 which encodes SEQ. ID NO. 1.

7. A substantially isolated nucleic acid polymer which encodes a protein which exhibits cytokinin oxidizing activity, wherein the nucleic acid polymer is selected from the group consisting of:

- 5 (a) SEQ. ID NO. 2,

(b) SEQ. ID NO. 3,

(c) a nucleic acid polymer with a sequence described by SEQ. ID NO. 10,

5 (d) a nucleic acid polymer whose sequence contains a sequence selected from the group consisting of SEQ. ID NO. 2, SEQ. ID NO. 3, or a nucleic acid polymer with a sequence described by SEQ. ID NO. 10.

10 (e) a nucleic acid polymer which contains at least one 60 base pair portion, which encodes an amino acid sequence which confers cytokinin oxidizing activity upon the encoded protein, of a nucleotide sequence selected from the group consisting of SEQ. ID NO. 2, SEQ. ID NO. 3, and a nucleic acid polymer with a sequence described by SEQ. ID NO. 10,

15 (f) a nucleic acid polymer which encodes a protein including an amino acid sequence with at least about 65% sequence identity to SEQ. ID NO. 1, the remainder of amino acid residues being conservatively substituted,

20 (g) a nucleic acid polymer which hybridizes to a nucleotide sequence selected from the group consisting of SEQ. ID NO. 2, SEQ. ID NO. 3, and a nucleic acid polymer with a sequence described by SEQ. ID NO. 10, under a wash stringency equivalent to 0.5X to 2X SSC buffer, 0.1% SDS, at 55-65°C, and

25 (h) a nucleic acid polymer having a nucleotide sequence complementary to any of the nucleic acid sequences (a) - (g).

8. The substantially isolated nucleic acid polymer of claim 7 wherein the polymer contains a sequence described by SEQ. ID NO. 10 or a sequence complementary to a sequence described by SEQ. ID NO. 10.

9. The substantially isolated nucleic acid polymer of claim 7 wherein the polymer is SEQ. ID No. 2 or its complementary sequence.

10. The substantially isolated nucleic acid polymer of claim 7 wherein the polymer is SEQ. ID No. 3 or its complementary sequence.

11. A host cell transformed with a vector containing a deoxyribonucleic acid polymer which encodes the protein of claim 1.

12. The host cell of claim 11, wherein the host cell is a plant cell.

13. The host cell of claim 12, wherein the host plant cell is selected from the group consisting of:

- (a) *Nicotiana tabacum*,
- (b) *Arabidopsis thaliana*,
- (c) *Zea mays*
- (d) *Brassica spp*
- (e) *Oryza sativa*

5

14. The host cell of claim 11 wherein the host cell is selected from the group consisting of:

- (a) *Pichia pastoris*
- (b) *Escherichia coli*

15. A plant regenerated from the host plant cell in claim 12.

16. A plant product from the regenerated plant of claim 15.

17. The plant product of claim 16, wherein said plant product is selected from the group consisting of seeds, leaves, stem cultures, rhizomes, and bulbs.

18. A method for producing the protein of claim 1 in a host cell, essentially comprising

- (a) constructing a vector containing DNA encoding the protein of claim 1 and an expression regulatory sequence operational in the host cell;
- (b) transfecting a host cell suitable for protein production with the vector; and
- 5 (c) expressing the protein in the host cell.

19. The method of claim 18 wherein the vector is selected from the group consisting of:

- (a) a plasmid comprised of DNA,
- (b) *Agrobacterium tumefaciens*

20. The method of claim 18, wherein the host cell is selected from the group consisting of:

- (a) *Pichia pastoris*
- (b) *Escherichia coli*

21. The method of claim 18 wherein the host cell secretes said protein into a culture medium, and wherein the method further comprises purifying said protein from the culture medium.

22. The method of claim 18 further comprising the step of reconstituting the host cell into an organism.

23. The method of claim 18, wherein the host cell is a plant cell.

24. The method of claim 23, wherein the host plant cell is selected from the group consisting of:

- (a) *Nicotiana tabacum*,
- (b) *Arabidopsis thaliana*,
- 5 (c) *Zea mays*
- (d) *Brassica spp*
- (e) *Oryza sativa*

25. A method of moderating cytokinin-associated pathogenesis in a plant comprising transforming a plant cell with a nucleic acid polymer construct which contains:

- 5 (a) a nucleic acid polymer encoding the protein of claim 1, and
  - (b) an expression regulatory sequence operational in the host plant; and
- regenerating a plant from the transformed plant cell.

26. A method as set forth in claim 25 wherein the expression regulatory sequence is selected from the group consisting of constitutive promoters, inducible promoters, repressors, and enhancers.

27. A method of detecting cytokinin concentrations in a sample essentially comprising:

- (a) adding to a buffered sample known amounts of dichlorophenolindophenol and an excess IU activity
- 5 producing amount of the protein of claim 1;
- (b) measuring the net change in absorbance of light at 590 nm; and
- (c) comparing this net absorbance to a standard curve generated from known concentrations of cytokinin.

FIG. 1b





FIG. 1a

 $N^6$ -( $\Delta^2$ -isopentenyl)adenine

3-methyl-2-butenal

FIG. 2A  
VERIFICATION OF *ckx1* GENE INTRON  
LOCATIONS BY RT-PCR  
1 2 3 4 5 6



FIG. 2B  
VERIFICATION OF *ckx1* GENE INTRON  
LOCATIONS BY RT-PCR  
1 2 3 4 5 6 7




FIG.3




FIG. 4




FIG. 5




FIG. 6





FIG. 7








FIG. 11




FIG. 12




FIG. 13




FIG.14





FIG.15



## SEQUENCE LISTING

&lt;110&gt; Morris Ph.D., Roy O.

&lt;120&gt; A CYTOKININ OXIDASE

&lt;130&gt; UMO 1491

<140>  
<141>

5

<150> 06/054,268  
<151> 1997-07-30

&lt;160&gt; 20

&lt;170&gt; PatentIn Ver. 2.0

10

<210> 1  
<211> 534  
<212> PRT  
<213> Zea mays<400> 1  
Met Ala Val Val Tyr Tyr Leu Leu Leu Ala Gly Leu Ile Ala Cys Ser  
1 5 10 15

15

His Ala Leu Ala Ala Gly Thr Pro Ala Leu Gly Asp Arg Gly Arg  
20 25 30

5 Pro Trp Pro Ala Ser Leu Ala Ala Leu Ala Leu Asp Gly Lys Leu Arg  
 35 40 45  
 Thr Asp Ser Asn Ala Thr Ala Ala Ser Thr Asp Phe Gly Asn Ile  
 50 55 60  
 Thr Ser Ala Leu Pro Ala Ala Val Leu Tyr Pro Ser Ser Thr Gly Asp  
 65 70 75 80  
 Leu Val Ala Leu Leu Ser Ala Ala Asn Ser Thr Pro Gly Trp Pro Tyr  
 85 90 95  
 Thr Ile Ala Phe Arg Gly Arg Gly His Ser Leu Met Gly Gln Ala Phe  
 100 105 110  
 Ala Pro Gly Gly Val Val Asn Met Ala Ser Leu Gly Asp Ala Ala  
 115 120 125  
 Ala Pro Pro Gly Ile Asn Val Ser Ala Asp Gly Arg Tyr Val Asp Ala  
 130 135 140  
 Gly Gly Glu Gln Val Trp Ile Asp Val Leu Arg Ala Ser Leu Ala Arg  
 145 150 155 160  
 Gly Val Ala Pro Arg Ser Trp Asn Asp Tyr Leu Tyr Leu Thr Val Gly  
 165 170 175  
 Gly Thr Leu Ser Asn Ala Gly Ile Ser Gly Gln Ala Phe Arg His Gly  
 180 185 190  
 Pro Gln Ile Ser Asn Val Leu Glu Met Asp Val Ile Thr Gly His Gly  
 195 200 205

"

Glu Met Val Thr Cys Ser Lys Gln Leu Asn Ala Asp Leu Phe Asp Ala  
 210 215 220  
 Val Leu Gly Gly Leu Gly Gln Phe Gly Val Ile Thr Arg Ala Arg Ile  
 225 230 235  
 Ala Val Glu Pro Ala Pro Ala Arg Ala Arg Trp Val Arg Phe Val Tyr  
 245 250 255  
 Thr Asp Phe Ala Ala Phe Ser Ala Asp Gln Glu Arg Leu Thr Ala Pro  
 260 265 270  
 Arg Pro Gly Gly Gly Ala Ser Phe Gly Pro Met Ser Tyr Val Glu  
 275 280 285  
 Gly Ser Val Phe Val Asn Gln Ser Leu Ala Thr Asp Leu Ala Asn Thr  
 290 295 300  
 Gly Phe Phe Thr Asp Ala Asp Val Ala Arg Ile Val Ala Leu Ala Gly  
 305 310 315  
 Glu Arg Asn Ala Thr Thr Val Tyr Ser Ile Glu Ala Thr Leu Asn Tyr  
 325 330 335  
 Asp Asn Ala Thr Ala Ala Ala Val Asp Gln Glu Leu Ala Ser  
 340 345 350

5

10

15

"

Val Leu Gly Thr Leu Ser Tyr Val Glu Gly Phe Ala Phe Gln Arg Asp  
 355 360 365  
 Val Ala Tyr Ala Ala Phe Leu Asp Arg Val His Gly Glu Glu Val Ala  
 370 375 380  
 Leu Asn Lys Leu Gly Leu Trp Arg Val Pro His Pro Trp Leu Asn Met  
 385 390 395 400  
 Phe Val Pro Arg Ser Arg Ile Ala Asp Phe Asp Arg Gly Val Phe Lys  
 405 410 415  
 Gly Ile Leu Gln Gly Thr Asp Ile Val Gly Pro Leu Ile Val Tyr Pro  
 420 425 430  
 Leu Asn Lys Ser Met Trp Asp Asp Gly Met Ser Ala Ala Thr Pro Ser  
 435 440 445  
 Glu Asp Val Phe Tyr Ala Val Ser Leu Leu Phe Ser Ser Val Ala Pro  
 450 455 460  
 Asn Asp Leu Ala Arg Leu Gln Glu Gln Asn Arg Arg Ile Leu Arg Phe  
 465 470 475 480  
 Cys Asp Leu Ala Gly Ile Gln Tyr Lys Thr Tyr Leu Ala Arg His Thr  
 485 490 495  
 Asp Arg Ser Asp Trp Val Arg His Phe Gly Ala Ala Lys Trp Asn Arg  
 500 505 510  
 Phe Val Glu Met Lys Asn Lys Tyr Asp Pro Lys Arg Leu Leu Ser Pro  
 515 520 525

5  
 10  
 15  
 20  
 "

Gly Gln Asp Ile Phe Asn  
530

<210> 2  
<211> 6733  
<212> DNA  
<213> Zea mays  
  
<220>  
<221> gene  
<222> (1)..(6733)  
<223> genomic sequence for a cytokinin oxidase from Zea  
mays  
  
<220>  
<221> CDS  
<222> (1497)..(2111)  
  
<220>  
<221> CDS  
<222> (2524)..(3216)  
  
<220>  
<221> CDS  
<222> (3311)..(3607)  
  
<220>  
<221> unsure  
<222> (5697)

5

10

15

20

25

<400> 2  
ctcgagactc cgacgagagg aggctgcgca tgcttgagtc atatcttga aaaaaaact 60  
gtaacttaaa gstatgatcta tatatggatt atttggatgg gatgtcattt tcgtatcacc 120  
aaccaaaatt acagttttggt cgtgcgtaga aattctacct actagctgaa acaacggctg 180  
ctatgtataa ctactgggtac tggaaagaat attagtcatt gactcaaaat tagaatgcat 240  
gtgtaagtca tgcgtgctaa tttgtttctat cagcattcgg cgaattccga agtccgtacg 300  
tgttgttcgt ggaggagagg aaaacatcag aaatgacaaa actagacggt gtgtgcttct 360  
acactgaatt cctcaacatt tgttttactt ttactagaga atggcatcaa atggaaaacc 420  
gctggaaaaa aaacaacaaa acaattggac ccaaatatg tatacagacg ctagctatag 480  
ccagccacac tgaagttgac atgcggcagc tagctagcca ctttctctga aacactaaca 540  
tttgtacctt ggtcgtgtaa gtgtagtagt aacgtatgtt gacgcgactt accgaacaaa 600  
aatataattg tcccaatcaa gctagggaagc attgttttgtt tccaaaatgt tgccatttgc 660  
ttaatcaatc ctatatattaat tcatggctgt taaggtagaga taaagcgaca agaaatctat 720  
atatatgtat ataagatccc gaaggctagc gacatttttg atagcaaaat atgagaagt 780  
tggcagattg ttctggtagc aaatcaaata atatggccag aataatcgtg gctagcttga 840  
ttaaaccttc atcagcttgg tgtatttttg aagtcgacca accagctggg ggtcgtcgtg 900  
cgtagtacca aaattacagc ctgctttcct tcgtcctgta cgtaatgcag tacagctgtc 960

tagtagagac cattttgagg aggcacacac acattaagtg ataacataaa agacggcctg 1020  
 attttatttc ataaccaaac gatattggtca acacacacct atagctacca aatttgtaca 1080  
 actatttagt gcgaaaacta ttctattctc aagaattgat cgcttatatt tattattaca 1140  
 gctttttaaa tgtataaata cgctatatattg catggcaaca gggggtaata attaggcagg 1200  
 actatatata taatagtttt ttctttctct gtaaattctt gggaggatgg taaagttggt 1260  
 aactaggcac ctacttgcg cgcataatttt tctgtgtgca aacagaataa aactagacgg 1320  
 gatgcagaat attttttcc ttggaaagca gctcatcttt gtgttcgagt aattgaagaa 1380  
 gtatgtaatc gcactacacc tacacctata tatatacggg gtgcaatcac ctagttacca 1440  
 aacactcaca cataacgtat agctctctct ctctcccggtg aacgacgacg tcgcta atg 1499  
 Met 1

gcg gtg gtt tat tac ctg ctg gcc ggg ctg atc gcc tgc tct cat 1547  
 Ala Val Val Tyr Tyr Leu Leu Ala Gly Leu Ile Ala Cys Ser His 15

gca cta gcg gca ggc acg cct gcg ctc gga gac gat cgc ggc cgt ccc 1595  
 Ala Leu Ala Ala Gly Thr Pro Ala Leu Leu Gly Asp Arg Gly Arg Pro 30

tgg cca gcc tcc ctc gcc gcg ctg gcc ttg gac ggc aag ctc cgg acc 1643  
 Trp Pro Ala Ser Leu Ala Ala Leu Ala Leu Asp Gly Lys Leu Arg Thr 45

5                   gac agc aac gcg acg gcg gcc gcg gac ttc ggc aac atc acg                   1691  
                   Asp Ser Asn Ala Thr Ala Ala Ser Thr Asp Phe Gly Asn Ile Thr           65  
                   50                   55                   60  
                   tcg gcg ctc ccg gcg gcg gtc ctg tac ccg tcg tcc acg ggc gac ctg           1739  
                   Ser Ala Leu Pro Ala Ala Val Leu Tyr Pro Ser Ser Thr Gly Asp Leu       80  
                   70                   75                   80  
                   gtg gcg ctg ctg agc gcg gcc aac tcc acc ccg ggg tgg ccc tac acc           1787  
                   Val Ala Leu Leu Ser Ala Ala Asn Ser Thr Pro Gly Trp Pro Tyr Thr       95  
                   85                   90  
                   atc gcg ttc cgc gcc gcg cgc cac tcc ctc atg ggc cag gcc ttc gcc           1835  
                   Ile Ala Phe Arg Gly Arg Gly His Ser Leu Met Gly Gln Ala Phe Ala       110  
                   100                   105  
                   ccc ggc ggc gtc gtc aac atg gcg tcc ctg ggc gac gcc gcc gcg           1883  
                   Pro Gly Gly Val Val Val Asn Met Ala Ser Leu Gly Asp Ala Ala Ala       125  
                   115                   120  
                   ccg ccg cgc atc aac gtg tcc gcg gac ggc cgc tac gtg gac gcc gcc ggc           1931  
                   Pro Pro Arg Ile Asn Val Ser Ala Asp Gly Arg Tyr Val Asp Ala Gly       145  
                   130                   135  
                   ggc gag cag gtg tgg atc gac gtg ttg cgc gcg tcg ctg gcg cgc ggc           1979  
                   Gly Glu Gln Val Trp Ile Asp Val Leu Arg Ala Ser Leu Ala Arg Gly       160  
                   150                   155  
                   gtg gcg ccg cgc tcc tgg aac gac tac ctc tac acc gtc ggc ggc           2027  
                   Val Ala Pro Arg Ser Trp Asn Asp Tyr Leu Tyr Leu Thr Val Gly Gly       175  
                   165                   170

5  
 10  
 15  
 20

acg ctg tcc aac gca ggc atc agc ggc cag gcg ttc cgc cac ggc cca 2075  
 Thr Leu Ser Asn Ala Gly Ile Ser Gly Gln Ala Phe Arg His Gly Pro  
 180 185 190  
 cag ata tct aac gtg ctg gag atg gac gtt atc acc ggtacgtgtg 2121  
 Gln Ile Ser Asn Val Leu Glu Met Asp Val Ile Thr  
 195 200 205  
 cactactac tactttttcc ctcccttgca caagtgaca accacaccac agcaagcgag 2181  
 caaaagcttg tttttttttt acgtgccagt acacctgcat cgacttctgt tgcttgccac 2241  
 ggggcaaac cgtgttcaat cagccggatt gaaattcggt acctacattg cgaatcatat 2301  
 atttatttt ttagtattat tagtggtgca tggtggttaa tgtccgcgct gcaccggccg 2361  
 gccgccgcc cggccggcga gggcgggcga cgtctttaat aactagtcac aaatcagcat 2421  
 gcatgctggc tctgcagct ggtgcgttga cattgtgcct ttgttcgttt cggctaatag 2481  
 aattatatg ctgggggtgtt gactttgtgg tgategaacg ca ggc cat ggg gag 2535  
 Met Val Thr Cys Ser Lys Gln Leu Asn Ala Asp Leu Phe Asp Ala Val  
 210 215 220 225  
 atg gtg acg tgc tcc aag cag ctg aac gcg gac ctg ttc gac gcc gtc 2583  
 ctg ggc ggg ctg ggg cag ttc gga gtg atc acc cgg gcc cgg atc gcg 2631  
 Leu Gly Gly Leu Gly Gln Phe Gly Val Ile Thr Arg Ala Arg Ile Ala  
 230 235 240 245

	2679	gtg gag ccg gcg ccg gcg ccg gcg ggc ggc gac cag gcg ctg tgg gtg cgg ttc acc gcc ccg tac acc Val Glu Pro Ala Pro Ala Arg Ala Arg Trp Val Arg Phe Val Tyr Thr 255
5	2727	gac ttc gcg gcg ttc agc gcc gcg cag gcg ctg acc gcc ccg ccg cgg Asp Phe Ala Ala Phe Ser Ala Asp Gln Gln Arg Leu Thr Ala Pro Arg 270
	2775	ccc ggc ggc ggc ggc gcg tcg ttc ggc ccg atg agc tac gtg gaa ggg Pro Gly Gly Gly Gly Ala Ser Phe Phe Gly Pro Met Ser Tyr Val Glu Gly 285
10	2823	tcg gtg ttc gtg aac cag agc ctg gcg acc gac ctg gcg aac acg ggg Ser Val Phe Val Asn Gln Ser Leu Ala Thr Asp Leu Ala Asn Thr Gly 300
	2871	ttc ttc acc gac gcc gac gtc gcc gcg atc gtc gcg ctc gcc ggg gag Phe Phe Thr Asp Ala Asp Val Ala Arg Ile Val Ala Leu Ala Gly Glu 315
15	2919	cgg aac gcc acc acc gtg tac agc atc gag gcc acg ctc aac tac gac Arg Asn Ala Thr Thr Thr Val Tyr Ser Ile Glu Ala Thr Leu Asn Tyr Asp 330
	2967	aac gcc acg gcg gcg gcg gcg ggc ggc gac cag gag ctg gcc tcc gtg Asn Ala Thr Ala Ala Ala Val Ala Val Asp Gln Glu Leu Ala Ser Val 345
20	3015	ctg ggc acg ctg agc tac gtg gag ggg ttc gcg ttc cag cgc gac gtg Leu Gly Thr Leu Ser Tyr Val Glu Gly Phe Ala Phe Gln Arg Asp Val 360

3063	gcc tac gcg gcg ttc ctt gac gcg gtg cac ggc gag gag gcg gctg gcg ctc	385
	Ala Tyr Ala Ala Phe Leu Asp Arg Val His Gly Glu Glu Val Ala Leu	380
	375	
3111	aac aag ctg ggg ctg tgg cgg gcg gcg cac ccg tgg ctc aac atg ttc	400
	Asn Lys Leu Gly Leu Trp Arg Val Pro His Pro Trp Leu Asn Met Phe	395
	390	
3159	gtg ccg cgc tcg cgc atc gcc gac ttc gac cgc ggc gtg ttc aag ggc	415
	Val Pro Arg Ser Arg Ile Ala Asp Phe Asp Arg Gly Val Phe Lys Gly	410
	405	
3207	atc ctg cag ggc acc gac atc gtc ggc ccg ctc atc gtc tac ccc ctc	430
	Ile Leu Gln Gly Thr Asp Ile Val Gly Pro Leu Ile Val Tyr Pro Leu	425
	420	
3256	aac aaa tcc atgtacgtgt tgaatcgatc ggctagctag ctagctaggc	
	Asn Lys Ser	
	435	
3313	acgccccggc cggcctctga cgactcgacc ggtctttctg gggtttggtt ttcc atg	Met
3361	tgg gac gac ggc atg tcg gcg gcg acg ccg tct gag gac gctg ttc tac	450
	Trp Asp Asp Gly Met Ser Ala Ala Thr Pro Ser Glu Asp Val Phe Tyr	445
	440	
3409	gcg gtg tcg ctg ctc ttc tcg tcg gtg gcg ccc aac gac ctg gcg agg	465
	Ala Val Ser Leu Leu Phe Ser Ser Val Ala Pro Asn Asp Leu Ala Arg	460
	455	

3457  
ctg cag gag cag aac agg agg atc ctg cgc ttc tgc gac ctc gcc ggg  
Leu Gln Glu Gln Asn Arg Arg Ile Leu Arg Phe Cys Asp Leu Ala Gly  
475 480 485  
470

3505  
atc cag tac aag acc tac ctg gcg cgg cac acg gac cgc agt gac tgg  
Ile Gln Tyr Lys Thr Tyr Leu Ala Arg His Thr Asp Arg Ser Asp Trp  
490 495 500

3553  
gtc cgc cac ttc ggc gcc ggc aag tgg aat cgc ttc gtg gag atg aag  
Val Arg His Phe Gly Ala Ala Lys Trp Asn Arg Phe Val Glu Met Lys  
505 510 515

3601  
aac aag tac gac ccc aag agg ctg ctc tcc ccc ggc cag gac atc ttc  
Asn Lys Tyr Asp Pro Lys Arg Leu Leu Ser Pro Gly Gln Asp Ile Phe  
520 525 530

3657  
aac tga tgatgatagc ccatgcatgt tttagtttct tgggacaata atgtgaataa  
Asn

3717  
ttcccagcta gcggctatta attaatctgt atatatgattg atcactgcac tgatgtatgt  
caaggtcatt atttacgtta tggtaaaaaa aaaagtaccc tcgtcgtata ttctcgtttc  
atctgcccaa ccccagagt tttaaaaaatg cacgttgaga gcaaaatttc cacaccgttt  
atccatgtag tagattgggt gtatattgaa tccgtataaa tatatagcaa gacaaaattt  
attgcaatct gatccccgtac accataattc acatgaaatt agaataaccg aacaaaagga  
cttgtagta tgggtgccc ttgcactttt tttgtgtgcc tatttaggta tgatatagtt  
ctgcaaaaca ttatttgaca gtcaaatata tccctgtcgt cctaataggc gtccaaaatc

4077

5      aaacgtgtgt ttagccagcc cccatccaaa cagcccata tcctatttt ctttgccttc 4137  
ccttttaatt ttgtttttt ttctctcttc aaataataac tggctcctca ctagtcactt 4197  
aaaaaaaaag aagagtgtga aggagaagga aaaagacacc gtcacttgtg aaacaaaatta 4257  
aaaagtttct tggatcgagg cgcgtgatac tctctcacc tgcaagtttt gcttcgatct 4317  
cgcacggcac cgtccgtcat ttctacaaa atacgaaacc ttgttccgtg cttgttgatc 4377  
agacagctga cgtcgacaag ttcaggtaaa gggtcatttg cgcgaccagc aggtcggctg 4437  
gtgttgaact gattctttta aacttaatga taaccagtag agtagacacc attgattctt 4497  
ttaaacttta accgctagtc ttattcaaaa tatttatta aaatgtataa ttttaaatca 4557  
agcacagact acatcaagtg ataaaaaaa tcataataaa attaatgata acttattatt 4617  
gttttgaata agacgaacga tcaaagtttt aaaaataaac ggtgtcatat acggagggag 4677  
tactagctta gtaccccggt tgaccagaag tgcgggtgtc cttaattcga tcgttttctc 4737  
ttgtttcctc agcgttcacc acatgcactg cacttgcaag ccgtacacgt tgactagtgg 4797  
ggcaatgcta gcttgggtgag gttttttgat tgattcccag cttttagacg aggtcactcc 4857  
acaaaaaaag ctaagcaacg gtgctgacct gaataataaa aggtgggtcgt agattattgg 4917  
ttatagccgt cgtcacaggt cacacacacg caacaagcct atagcaggat tacaccgatt 4977  
atcatcggga aatagaaaaa agggaaatga acgaagtaac caggagacag acgacgagaa 5037  
agtattgtct atccttaaga ccccttgag caagctacac ggataattag gctcccatcg 5097

10      15      20      25      "

30

gacgtcattt ctggacacgt acgctattgt gatttgaag ggtgtttggt ttctaaatac 5157  
taatttttag ccactcttta ttattatatt ctagtcacta aattatcaaa tacgaaaaatt 5217  
aaaaatagatt tttaatttta agtatttttgt aattttatgta ctagaatgga ataaaaatggt 5277  
gtgattaaaa attagtcctt aaaaatcaaa tgtcattccc cctttattag aagacctctt 5337  
gagcaagcta agctgcacgt attattagc ctcaacatgc ctatgtgtgt ataacaggac 5397  
aaccaggccc cccaaaactc agttgtacgt atgtgtaaca tgtactgtga ggacaatcga 5457  
tctcaaacgt cagctgtagg tacgtatgaa tgtatatatg tgtatttact cctcctgttt 5517  
taaatcagtt gtcgcgatgg tatttatgtc catcaaagtt tgttagagta gactagcttt 5577  
ttagaaaaatg ttagtcacat ttatatcctc aaataaaactt acttataaaa ataaattcaa 5637  
tgatctattt aataaatataa cgatactaata catgtaatat gaataataaat attttcttgn 5697  
acatatgtga tgaagttaa aatagtttagt ttttaaaaaa ataaaaaacac cgactatttt 5757  
aaaatagata gagtacatat acaagaaaat gagaaattat tactaatgtt tacaacgtat 5817  
aataaataat atataaaatta tgaaatatat ttttataata aatctgttta gagaagataa 5877  
tatttttctac gactccttgt tcaaattctag aatgacatta tttttgggac ggaggcagtt 5937  
cgtactacgt acgtaacatg tattcctaac acggacaatc aaaacatggg aacttcactc 5997  
catttgtaaa gccggggggcg gaatcggtgg amgtgtttgt ttcttggcgc acggtgaytc 6057  
gatgaggatt cgtgcaaacc ttaamaggct tggatcatctg tctgcattca ytcacacgg 6117

5

10

15

20

"

ccatacgcac gtgtttttct ggatgggcat gaacacggac acggcacgca tctcaagtca 6177  
gggcagttaa agggagacgc gtgggaacat gataccaggg ttgcatcttc acaaaattgg 6237  
ccgtatgtgc tacatcatca gggggagcag ccataattta catggaaacg aatacaatga 6297  
tggagcacgt cagtggctat tattcttcaa tgaggctaca gctacactat ttgatgcaca 6357  
tcagctactg cctcctagtc cttggggcgc cacctccaca ctttcagctc gccggagcag 6417  
tctccgggtga agagcagccc accagccgca agctcgatgg tcctaaacttc cttcgtggac 6477  
aacagtggtc caatcccgtc gaacctgaaa gatggtaacc aggcaaatgc agagctcctt 6537  
tcagtaagga accactgagc gagcggtttg tctatctgat gatgaaactg ataaaaaaaa 6597  
attatgcact cacgatggca agtcgagcaa gcggacgcga ttgctgttgt ggaggaggca 6657  
caagagaact ggcgtcttcc caactcggtg catgccgaag agtgtgcgca aaccctgcaa 6717  
cataatcgga aagctt 6733

<210> 3  
<211> 1605  
<212> DNA  
<213> Zea mays

<220>  
<221> CDS  
<222> (1)..(1605)

"

<400> 3  
 atg gcg gtg gtt tat tac ctg ctg ctg gcc ggg ctg atc gcc tgc tct 48  
 Met Ala Val Tyr Tyr 5  
 10  
 15  
 20  
 cat gca cta gcg gca ggc acg cct gcg ctg gga gac gat cgc ggc cgt 96  
 His Ala Leu Ala Ala Gly Thr Pro Ala Leu Gly Asp Arg Gly Arg  
 25  
 30  
 35  
 40  
 45  
 50  
 55  
 60  
 65  
 70  
 75  
 80  
 85  
 90  
 95  
 100  
 105  
 110  
 115  
 120  
 125  
 130  
 135  
 140  
 145  
 150  
 155  
 160  
 165  
 170  
 175  
 180  
 185  
 190  
 195  
 200  
 205  
 210  
 215  
 220  
 225  
 230  
 235  
 240  
 245  
 250  
 255  
 260  
 265  
 270  
 275  
 280  
 285  
 290  
 295  
 300  
 305  
 310  
 315  
 320  
 325  
 330  
 335  
 340  
 345  
 350  
 355  
 360  
 365  
 370  
 375  
 380  
 385  
 390  
 395  
 400

432

gac ccg ccg ccg atc aac gtg tcc gac gac gac gcc  
Ala Pro Pro Arg Ile Asn Val Ser Ala Asp Gly  
130 135 140

480  
 ggc ggc gag cag gtg tgg atc gac gtg ttg cgc ggc tcg ctg ggc cgc  
 Gly Gly Glu Gln Val Trp Ile Asp Val Leu Arg Ala Ser Leu Ala Arg  
 145 150 155 160

528  
 ggc gtg ggc ccg cgc tcc tgg aac gac tac ctc tac ctc acc gtc ggc  
 Gly Val Ala Pro Arg Ser Trp Asn Asp Tyr Leu Tyr Leu Thr Val Gly  
 165 170 175

576  
 ggc acg ctg tcc aac gca ggc atc agc ggc cag ggc ttc cgc cac ggc  
 Gly Thr Leu Ser Asn Ala Gly Ile Ser Gly Gln Ala Phe Arg His Gly  
 180 185 190

624  
 cca cag ata tct aac gtg ctg gag atg gac gtt atc acc ggc cat ggc  
 Pro Gln Ile Ser Asn Val Leu Glu Met Asp Val Ile Thr Gly His Gly  
 195 200 205

672  
 gag atg gtg acg tgc tcc aag cag ctg aac ggc gac ctg ttc gac gcc  
 Glu Met Val Thr Cys Ser Lys Gln Leu Asn Ala Asp Leu Phe Asp Ala  
 210 215 220

720  
 gtc ctg ggc ggc ctg ggc cag ttc gga gtg atc acc cgc gcc cgc atc  
 Val Leu Gly Gly Leu Gly Gln Phe Gly Val Ile Thr Arg Ala Arg Ile  
 225 230 235 240

768  
 ggc gtg gag ccg ggc ccg ggc cgc ggc ccg ggc ggc tgg gtg ttc gtg tac  
 Ala Val Glu Pro Ala Pro Ala Arg Ala Arg Trp Val Arg Phe Val Tyr  
 245 250 255

816  
 acc gac ttc ggc ggc ttc agc gcc gac cag gag cgc ctg acc gcc ccg  
 Thr Asp Phe Ala Ala Phe Ser Ala Asp Gln Glu Arg Leu Thr Ala Pro  
 260 265 270



1248  
 ttc gtg ccg cgc tcg cgc atc gcc gac ttc gac cgc arg gly val phe lys  
 phe val pro arg ser arg ile ala asp phe asp arg gly val phe lys  
 405 410 415  
 1296  
 ggc atc ctg cag ggc acc gac atc gtc ggc ccg ctc atc gtc tac ccc  
 gly ile leu gln gly thr asp ile val gly pro leu ile val tyr pro  
 420 425 430  
 1344  
 ctc aac aaa tcc atg tgg gac gac ggc atg tcg gcg gcg acg ccg tct  
 leu asn lys ser met trp asp ggc asp gly met ser ala ala thr pro ser  
 435 440 445  
 1392  
 gag gac gtg ttc tac gcg gtg tcg ctg ctc ttc tcg gtc gtc ggc ccc  
 glu asp val phe tyr ala val ser leu leu phe phe ser val ala pro  
 450 455 460  
 1440  
 aac gac ctg gcg agg ctg cag gag gln gln gln arg arg ile leu arg phe  
 asn asp leu ala arg leu 470 475 480  
 1488  
 tgc gac ctc gcc ggg atc cag tac aag acc acc tcg gcg cgc cac acg  
 cys asp leu ala gly ile gln tyr lys thr tyr leu ala arg his thr  
 485 490 495  
 1536  
 gac cgc agt gac tgg gtc cgc cac ttc ggc gcc gcc aag tgg aat cgc  
 asp arg ser asp trp val arg his phe gly ala ala lys trp asn arg  
 500 505 510  
 1584  
 ttc gtg gag atg aag aac aag tac gac ccc aag agg ctg ctc tcc ccc  
 phe val glu met lys asn lys tyr asp pro lys arg leu leu ser pro  
 515 520 525

1605

ggc cag gac atc ttc aac tga  
Gly Gln Asp Ile Phe Asn  
530

<210> 4  
<211> 21  
<212> PRT  
<213> Zea mays

<220>  
<221> VARIANT  
<222> (9)

<400> 4  
Tyr Val Glu Gly Ser Val Phe Val Xaa Gln Ser Leu Ala Thr Asp Leu  
1 5 10 15

Ala Asn Thr Gly Phe  
20

<210> 5  
<211> 15  
<212> PRT  
<213> Zea mays

<400> 5  
Gly Ile Leu Gln Gly Thr Asp Ile Val Gly Pro Leu Ile Val Tyr  
1 5 10 15

<210> 6  
<211> 23  
<212> DNA  
<213> Zea mays

<220>  
<221> modified\_base  
<222> (6)  
<223> i

<220>  
<221> modified\_base  
<222> (12)  
<223> i

<220>  
<221> modified\_base  
<222> (15)  
<223> i

<220>  
<221> modified\_base  
<222> (18)  
<223> i

<400> 6  
tacgtngayg gnwsngtntt cgt

<210> 7  
<211> 23  
<212> DNA  
<213> Zea mays

23

5

10

15

20

25

30

23

<220>  
<221> modified\_base  
<222> (3)  
<223> i

5

<220>  
<221> modified\_base  
<222> (9)  
<223> i

10

<220>  
<221> modified\_base  
<222> (12)  
<223> i

15

<220>  
<221> modified\_base  
<222> (18)  
<223> i

20

<220>  
<221> modified\_base  
<222> (21)  
<223> i

25

<400> 7  
acngayctng cnaacacngg ntt

<210> 8  
<211> 23  
<212> DNA  
<213> Zea mays

30

23

<220>  
<221> modified\_base  
<222> (3)  
<223> i

5

<220>  
<221> modified\_base  
<222> (9)  
<223> i

10

<220>  
<221> modified\_base  
<222> (15)  
<223> i

15

<220>  
<221> modified\_base  
<222> (18)  
<223> i

<400> 8  
ccntargang tccntgnct rta

20

<210> 9  
<211> 23  
<212> DNA  
<213> Zea mays

25

<220>  
<221> modified\_base  
<222> (6)  
<223> i

"

23

<220>  
<221> modified\_base  
<222> (9)  
<223> i

5

<220>  
<221> modified\_base  
<222> (12)  
<223> i

10

<220>  
<221> modified\_base  
<222> (15)  
<223> i

15

<220>  
<221> modified\_base  
<222> (21)  
<223> i

20

<400> 9  
tarcancng gngantarca nat

25

<210> 10  
<211> 1602  
<212> DNA  
<213> Zea mays

30

<220>  
<221> variation  
<222> (6)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (9)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (12)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (21)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (24)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (27)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (30)  
<223> a,g,c or t

30

<220>  
<221> variation  
<222> (33)

<223> a, g, c or t

<220>  
<221> variation  
<222> (36)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (42)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (48)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (54)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (57)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (60)  
<223> a, c, g or t

" 30

<220>  
<221> variation  
<222> (63)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (66)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (69)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (72)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (75)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (78)  
<223> a, g, c or t

30

"

<220>  
<221> variation  
<222> (81)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (90)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (93)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (96)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (99)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (105)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (108)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (111)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (114)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (117)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (120)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (123)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (126)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (129)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (135)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (141)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (144)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (147)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (153)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (159)  
<223> a, c, g or t

10

<220>  
<221> variation  
<222> (162)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (165)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (168)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (171)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (174)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (177)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (186)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (195)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (198)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (201)  
<223> a,g,c or t

5

10

15

20

25

"

<220>  
<221> variation  
<222> (204)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (207)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (210)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (213)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (216)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (219)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (225)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (228)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (231)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (234)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (237)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (243)  
<223> a,g,c or t

5

10

15

20

25

"

<220>  
<221> variation  
<222> (246)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (249)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (252)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (255)  
<223> a,c,g or t

20

<220>  
<221> variation  
<222> (258)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (261)  
<223> a,g,c or t

"

<220> variation  
<221> (264)  
<222> a, g, c or t

<220> variation  
<221> (270)  
<222> a, g, c or t

<220> variation  
<221> (273)  
<222> a, g, c or t

<220> variation  
<221> (276)  
<222> a, g, c or t

<220> variation  
<221> (279)  
<222> a, g, c or t

<220> variation  
<221> (285)  
<222> a, g, c or t

5

10

15

20

25

"

<220>  
<221> variation  
<222> (291)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (297)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (303)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (306)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (309)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (312)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (318)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (321)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (327)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (333)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (339)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (342)  
<223> a,g,c or t

25

"

<220>  
<221> variation  
<222> (345)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (348)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (351)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (354)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (357)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (366)  
<223> a,c,g or t

"

<220>  
<221> variation  
<222> (369)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (372)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (375)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (381)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (384)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (387)  
<223> a, g, c or t

5

10

15

20

25

"

<220>  
<221> variation  
<222> (390)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (393)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (396)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (405)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (408)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (411)  
<223> a, g, c or t

"

<220> variation  
<221> (417)  
<222> a, g, c or t

<220> variation  
<221> (420)  
<222> a, g, c or t

<220> variation  
<221> (426)  
<222> a, g, c or t

<220> variation  
<221> (432)  
<222> a, g, c or t

<220> variation  
<221> (435)  
<222> a, g, c or t

<220> variation  
<221> (438)  
<222> a, g, c or t

5

10

15

20

25

<220>  
<221> variation  
<222> (447)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (459)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (462)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (465)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (468)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (471)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (474)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (477)  
<223> a,c,g or t

<220>  
<221> variation  
<222> (480)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (483)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (486)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (489)  
<223> a,g,c or t

5

10

15

20

"

<220>  
<221> variation  
<222> (492)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (495)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (498)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (513)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (519)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (522)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (525)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (528)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (531)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (534)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (537)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (540)  
<223> a,g,c or t

5

10

15

20

"

<220>  
<221> variation  
<222> (546)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (549)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (555)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (558)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (564)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (570)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (576)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (579)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (588)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (594)  
<223> a,c,g or t

20

<220>  
<221> variation  
<222> (597)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (609)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (615)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (618)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (624)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (633)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (637)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (642)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (651)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (657)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (663)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (672)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (675)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (678)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (681)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (684)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (687)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (690)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (699)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (702)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (708)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (711)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (717)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (723)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (726)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (732)  
<223> a, c, g or t

"

<220>  
<221> variation  
<222> (735)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (738)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (741)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (744)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (747)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (750)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (756)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (759)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (765)  
<223> a,g,c or t

&lt;220&gt;

<221> variation  
<222> (771)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (780)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (783)  
<223> a,g,c or t

25

"

<220>  
<221> variation  
<222> (789)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (792)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (804)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (807)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (810)  
<223> a, g, c or t

25 "

<220>  
<221> variation  
<222> (813)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (816)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (819)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (822)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (825)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (828)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (831)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (834)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (837)  
<223> a, c, g or t

10

<220>  
<221> variation  
<222> (840)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (846)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (849)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (855)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (861)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (867)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (870)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (873)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (879)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (888)  
<223> a,g,c or t

5

10

15

20

"

<220>  
<221> variation  
<222> (891)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (894)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (897)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (903)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (906)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (912)  
<223> a, g, c or t

"

5  
    <220> variation  
    <221> (915)  
    <222> a, g, c or t  
    <223> a, g, c or t  
10  
    <220> variation  
    <221> (924)  
    <222> a, g, c or t  
    <223> a, g, c or t  
15  
    <220> variation  
    <221> (930)  
    <222> a, g, c or t  
    <223> a, g, c or t  
20  
    <220> variation  
    <221> (936)  
    <222> a, g, c or t  
    <223> a, g, c or t  
25  
    <220> variation  
    <221> (942)  
    <222> a, c, g or t  
    <223> a, c, g or t  
"

<220>  
<221> variation  
<222> (948)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (951)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (954)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (957)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (960)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (966)  
<223> a,g,c or t

5

10

15

20

25

"

<220>  
<221> variation  
<222> (972)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (975)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (978)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (987)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (996)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (999)  
<223> a, g, c or t

5

10

15

20

"

<220>  
<221> variation  
<222> (1002)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (1017)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (1020)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (1023)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (1026)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (1029)  
<223> a,g,c or t

25

"

<220>  
<221> variation  
<222> (1032)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1035)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1038)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1050)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1053)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1056)  
<223> a, g, c or t

5

10

15

20

25

"

<220>  
<221> variation  
<222> (1059)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (1062)  
<223> a, c, g or t

10

<220>  
<221> variation  
<222> (1065)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (1068)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (1071)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (1074)  
<223> a, g, c or t

"

5  
10  
15  
20  
25  
"

<220> variation  
<221> (1080)  
<222> a, g, c or t  
<223> a, g, c or t

<220> variation  
<221> (1086)  
<222> a, g, c or t  
<223> a, g, c or t

<220> variation  
<221> (1092)  
<222> a, g, c or t  
<223> a, g, c or t

<220> variation  
<221> (1101)  
<222> a, g, c or t  
<223> a, g, c or t

<220> variation  
<221> (1107)  
<222> a, g, c or t  
<223> a, g, c or t

<220> variation  
<221> (1110)  
<222> a, g, c or t  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1116)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (1119)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (1125)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (1131)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (1134)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (1140)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (1149)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (1152)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (1155)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (1164)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (1167)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (1170)  
<223> a,g,c or t

"

<220>  
<221> variation  
<222> (1176)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (1182)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (1188)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (1194)  
<223> a, c, g or t

20

<220>  
<221> variation  
<222> (1206)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (1209)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (1212)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1215)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1218)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1224)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1236)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1239)  
<223> a, g, c or t

5

10

15

20

"

5	<220> <221> variation <222> (1242) <223> a,g,c or t
10	<220> <221> variation <222> (1251) <223> a,g,c or t
15	<220> <221> variation <222> (1263) <223> a,g,c or t
20	<220> <221> variation <222> (1266) <223> a,g,c or t
25	<220> <221> variation <222> (1275) <223> a,g,c or t
"	

<220>  
<221> variation  
<222> (1278)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1281)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1284)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1290)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1296)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1299)  
<223> a, g, c or t

5

10

15

20

"

<220>  
<221> variation  
<222> (1308)  
<223> a,g,c or t

5

<220>  
<221> variation  
<222> (1323)  
<223> a,g,c or t

10

<220>  
<221> variation  
<222> (1329)  
<223> a,g,c or t

15

<220>  
<221> variation  
<222> (1332)  
<223> a,g,c or t

20

<220>  
<221> variation  
<222> (1335)  
<223> a,g,c or t

25

<220>  
<221> variation  
<222> (1338)  
<223> a,c,g or t

"

<220>  
<221> variation  
<222> (1341)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (1344)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (1353)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (1362)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (1365)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (1368)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1371)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (1374)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (1380)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (1383)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (1386)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (1389)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (1392)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (1401)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (1404)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (1410)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (1425)  
<223> a,g,c or t

<220>  
<221> variation  
<222> (1428)  
<223> a,g,c or t

5

10

15

20

<220>  
<221> variation  
<222> (1434)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (1437)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (1449)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (1452)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (1455)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (1470)  
<223> a, g, c or t

"

<220> variation  
<221> (1476)  
<222> a, g, c or t

5

<220> variation  
<221> (1479)  
<222> a, c, g or t

10

<220> variation  
<221> (1482)  
<222> a, g, c or t

15

<220> variation  
<221> (1488)  
<222> a, g, c or t

20

<220> variation  
<221> (1494)  
<222> a, g, c or t

25

<220> variation  
<221> (1497)  
<222> a, g, c or t

"

<220>  
<221> variation  
<222> (1506)  
<223> a, g, c or t

5

<220>  
<221> variation  
<222> (1509)  
<223> a, g, c or t

10

<220>  
<221> variation  
<222> (1518)  
<223> a, g, c or t

15

<220>  
<221> variation  
<222> (1521)  
<223> a, g, c or t

20

<220>  
<221> variation  
<222> (1524)  
<223> a, g, c or t

25

<220>  
<221> variation  
<222> (1536)  
<223> a, g, c or t

"

<220>  
<221> variation  
<222> (1542)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1566)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1572)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1575)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1578)  
<223> a, g, c or t

<220>  
<221> variation  
<222> (1581)  
<223> a, g, c or t

5

10

15

20

25

"

<220>  
 <221> variation  
 <222> (1584)  
 <223> a, g, c or t

5

<220>  
 <221> variation  
 <222> (1587)  
 <223> a, g, c or t

10

<400> 10  
 atggcngtng tntaytayyt nytnytnngcn ggnytnathg cntgywsnca ygcnytnngcn 60  
 gcnggnacnc cngcnytnngg ngaygaymgn ggnmgncnt gccngcncws nytnngcngcn 120  
 ytnngcnytnng ayggnaaryt nmgnacngay wsnaaygcna cngcngcngc nwsnacngay 180  
 ttyggnaaya thacnwsngc nytnccngcn gcngtnytnnt ayccnwsnws nacngngngay 240  
 ytngtngcny tnytnwsngc ngcnaaywsn acncnggnt gccntayac nathgcntty 300  
 mngngnmng gncaywsnyt natggngcar gcnttygcnc cnggnggngt ngtnngtnaay 360  
 atggcnwsny tnggngaygc ngcngcncn ccmgnatha aygtnwsngc ngayggnmgn 420  
 taygtngayg cnggnggnga rcargtntgg athgaygtny tnmngcncws nytnngcnmgn 480  
 ggngtngcnc cnmgnwsntg gaaygaytay ytntayytna cngtnggngg nacnytnwsn 540  
 aaygcnggna thwsnggna rgcnttymgn cayggncnc arathwsnaa ygtnytnngar 600  
 atggaygtna thacnggna yggngaratg gtnacntgyw snaarcaryt naaygcngay 660

15

20

25

"

ytnttygayg cngtnytnng nggnytnngn carttyggng tnathacnmg ngcnmgcnath 720  
gcngtn Garc cngcncngc nmngncnmgn tgggtmngnt tygtntayac ngayttygc 780  
gcnttywsng cngaycarga rmgnytacn gcncncmgnc cngngggngg ngngncnwsn 840  
ttyggncna tgsntaygt ngarggnwsn gtnttygtna aycarwsnyt ngcnacngay 900  
ytnGcnaaya cnggnttytt yacngaygn gaygtngcnm gnathgtngc nytnGcnggn 960  
garmGnaayg cnacnacngt ntaywsnath gargcnacny tnaaytayga yaayGcnacn 1020  
gcngcngcng cngcngtnga ycargarytn gcnwsngtny tngGnacny nwsntaygt 1080  
garggnttyg cnttycarmg ngaygtngcn taygcngcnt tyytnGaymg ngtncaaygn 1140  
gargargtn cnytnaayaa rytnggnytn tggmgngtnc cncayccntg gytnaayatg 1200  
ttygtncnmm gnwsnmgnat hgcngaytty gaymgngng tnttyaargg nathytnear 1260  
ggnacngaya thgtnggncc nytnathgt 1320  
ggnatgwsng cngcnacncc nwsngargay gtnttytayg cngtnwsnyt nytnnttywsn 1380  
wsngtnGcnc cnaaygayt ngcnmgnytn cargarcara aymgnmgnt hytnmgntty 1440  
tgygayytng cnggnathca rtayaaraen tayytnGcnm gncayacnga ymgnwsngay 1500  
tgggtnmGnc aytytyggngc ngcnaartgg aaymgnttyg tngaratgaa raayaartay 1560  
gayccnaarm gnytnytnws nccnggnear gayathttya ay 1602

5  
    <210> 11  
    <211> 32  
    <212> DNA  
    <213> Zea mays  
    <400> 11  
tggaattcc atgggagat ggtgacgtgc tc  
32

10  
    <210> 12  
    <211> 33  
    <212> DNA  
    <213> Zea mays  
    <400> 12  
gccgtccac atggatttgt tgaggggta gac  
33

15  
    <210> 13  
    <211> 37  
    <212> DNA  
    <213> Zea mays  
    <400> 13  
ctcaacaaat ccatgtgga cgacggcatg tcggcgg  
37

20  
    <210> 14  
    <211> 39  
    <212> DNA  
    <213> Zea mays  
25

39

<400> 14  
gcggtctaga tctaactaaa acatgcatgg gctatcatc

33

<210> 15  
<211> 33  
<212> DNA  
<213> Zea mays

33

<400> 15  
atgggaattc catggggaga tggtagcgtg ctc

39

<210> 16  
<211> 39  
<212> DNA  
<213> Zea mays

39

<400> 16  
gcggtctaga tctaactaaa acatgcatgg gctatcatc

17

<210> 17  
<211> 17  
<212> DNA  
<213> Zea mays

<400> 17  
ccggttttgg taccggg

20

<210> 18  
<211> 17

"

<212> DNA  
<213> Zea mays

<400> 18  
catgaccggt accaaaa

17

<210> 19  
<211> 22  
<212> DNA  
<213> Zea mays

5

<400> 19  
gacaccattc caagcatacc cc

22

10

<210> 20  
<211> 22  
<212> DNA  
<213> Zea mays

<400> 20  
gttctcacta gaaaaatgcc cc

22

15

"

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/15844

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/53 C12N15/82 C12N15/81 C12N9/06 C12N1/21  
C12N1/19 C12N5/10 A01H5/00 C12Q1/26

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 119, no. 21, 22 November 1993 Columbus, Ohio, US; abstract no. 221752, BURCH, L. R. ET AL: "Cytokinin oxidase and the degradative metabolism of cytokinins" XP002086612 see abstract & PHYSIOL. BIOCHEM. CYTOKININS PLANTS, SYMP. (1992), MEETING DATE 1990, 29-32. EDITOR(S): KAMINEK, MIROSLAV;MOK, DAVID W. S.; ZAZIMALOVA, EVA. PUBLISHER: SPB ACAD. PUBL., THE HAGUE, NETH. CODEN: 59KXA9, --- -/--	1-4

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*S\* document member of the same patent family

Date of the actual completion of the international search

3 December 1998

Date of mailing of the international search report

18.01.99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

MADDOX, A

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/15844

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCHREIBER B M N ET AL: "Polyclonal antibodies to maize seedling cytokinin oxidase" ANNUAL MEETING OF THE AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, CHARLOTTE, NORTH CAROLINA, USA, JULY 29-AUGUST 2, 1995. PLANT PHYSIOLOGY (ROCKVILLE) 108 (2 SUPPL.). 1995. 80., XP002086604	1-4
A	see the whole document ---	5-24
X	BURCH, LINDSAY R. ET AL: "The purification of cytokinin oxidase from Zea mays kernal" PHYTOCHEMISTRY (1989), 28(5), 1313-19, XP002086605	1-4
Y	see the whole document ---	5-24
Y	MEILAN R ET AL: "Cloning the cytokinin oxidase gene." ANNUAL MEETING OF THE AMERICAN SOCIETY OF PLANT PHYSIOLOGISTS, PORTLAND, OREGON, USA, JULY 30-AUGUST 3, 1994. PLANT PHYSIOLOGY (ROCKVILLE) 105 (1 SUPPL.). 1994. 68., XP002086606 see the whole document ---	5-24
Y	MIROSLAV KAMINEK: "PROGRESS IN CYTOKININ RESEARCH" TRENDS IN BIOTECHNOLOGY, vol. 10, no. 5, 1 May 1992, pages 159-164, XP000272384 see page 162 ---	5-24
P,X	MORRIS, R.O, ET AL.: "Zea mays cytokinin oxidase (ckx1) gene, complete cds" EMBL SEQUENCE ACCESSION NO. AF044603, 27 July 1998, XP002086607 see the whole document ---	1-11
A	WANG, J. ET AL: "Cytokinin oxidase purification by affinity chromatography and activation by caffeic acid" PLANT SCI. (SHANNON, IREL.) (1995), 112(2), 161-6, XP002086608 see the whole document ---	1-24
A	MOTKYA, V., ET AL.: "Changes in cytokinin content and cytokinin oxidase activity in response to derepression of ipt gene transcripion in transgenic tobacco calli and plants" PLANT PHYSIOLOGY, vol. 112, 1996, pages 1035-1043, XP002086609 see the whole document ---	1-24

-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/15844

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CRESPI, M., ET AL.: "Hypothetical 47.9 kD oxidoreductase in fasciation locus(ORF5)"            SWISSPROT ACCESSION NO .P46377,            1 November 1995, XP002086610            see the whole document</p>	1-4
A	<p>---            CRESPI., M., ET AL.: "R. fascians (D188) genes for P450 cytochrome, isopentenyltransferase and ferridoxine"            EMBL ACCESSION NO. Z29635,            15 February 1994, XP002086611            see the whole document</p> <p>-----</p>	5-10



## REFERENCES

- WO0105985. Method to modulate the expression of genes inducing the parthenocarpic trait in plants.
- Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., and Watson, J. D. (1994).  
5 "Molecular Biology of the Cell." Garland Publishing Inc.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. & Lipman, D.J. (1997) "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs." *Nucl. Acids Res.* **25**, 3389-3402.
- Armstrong, D.J. (1994) in *Cytokinins: Chemistry, Activity and Functions*, eds.  
10 Mok. D.W.S & Mok, M.C. (CRC Boca Raton, FL), pp. 139-154.
- An, G., Watson, B. D., Stachel, S., Gordon, M. P., and Nester, E. W. (1985). New cloning vehicles for transformation of higher plants. *EMBO J.* **4**, 277-284.
- Armstrong, C. L., Petersen, W. P., Buchholz, W. G., Bowen, B. A., and Sulc, S. L. (1990). Factors affecting PEG-mediated stable transformation of maize  
15 protoplasts. *Plant Cell Reports* **9**, 335-339.
- Banerjee, A., Pramanik, A., Bhattacharjya, S., and Balaram, P. (1996). Omega amino acids in peptide design: incorporation into helices. *Biopolymers* **39**, 769-777.
- Baron, M. H. and Baltimore, D. (1982). Antibodies against the chemically  
20 synthesized genome-linked protein of poliovirus react with native virus-specific proteins. *Cell* **28**, 395-404.
- Bartel, P. L. and Fields, S. (1997). "The Yeast Two-Hybrid System." Oxford University Press.
- Benkirane, N., Guichard, G., Briand, J. P., and Muller, S. (1996). Exploration of  
25 requirements for peptidomimetic immune recognition. Antigenic and immunogenic properties of reduced peptide bond pseudopeptide analogues of a histone hexapeptide. *J. Biol. Chem.* **271**, 33218-33224.



- Berry, A. and Brenner, S. E. (1994). A prototype computer system for de novo protein design. *Biochem.Soc.Trans.* **22**, 1033-1036.
- Christou, P., McCabe, D. E., and Swain, W. F. (1988). Stable transformation of soybean callus by DNA-coated gold particles. *Plant Physiol.* **87**, 671-674.
- 5 Crossway, A., Oakes, J. V., Irvine, J. M., Ward, B., Knauf, V. C., and Shewmaker, C. K. (1986). Integration of foreign DNA following microinjection of tobacco mesophyll protoplasts. *Mol.Gen.Genet.* **202**, 179-185.
- Dale, E. C. and Ow, D. W. (1990). Intra- and intermolecular site-specific recombination in plant cells mediated by bacteriophage P1 recombinase. *Gene* **91**,  
10 79-85.
- Dodds, J. H. (1985). "Plant genetic engineering." Cambridge University Press.
- Doerner, P., Jorgensen, J. E., You, R., Steppuhn, J., and Lamb, C. (1996). Control of root growth and development by cyclin expression. *Nature* **380**, 520-523.
- Dorner, B., Husar, G. M., Ostresh, J. M., and Houghten, R. A. (1996). The  
15 synthesis of peptidomimetic combinatorial libraries through successive amide alkylations. *Bioorg.Med.Chem.* **4**, 709-715.
- Ellis, J. G., Llewellyn, D. J., Dennis, E. S., and Peacock, W. J. (1987). Maize Adh-1 promoter sequences control anaerobic regulation: addition of upstream promoter elements from constitutive genes is necessary for expression in tobacco.  
20 *EMBO J.* **6**, 11-16.
- Faiss, M., Zalubilová, J., Strnad, M., Schmülling, T. (1997). Conditional transgenic expression of the ipt gene indicates a function for cytokinins in paracrine signaling in whole tobacco plants. *Plant J.* **12**, 401-415.
- Fassina, G. and Melli, M. (1994). Identification of interactive sites of proteins and  
25 protein receptors by computer-assisted searches for complementary peptide sequences. *Immunomethods.* **5**, 114-120.



- Fedoroff, N. V. and Smith, D. L. (1993). A versatile system for detecting transposition in Arabidopsis. *Plant J.* **3**, 273-289.
- Hanahan, D. (1983). Studies on transformation of Escherichia coli with plasmids. *J.Mol.Biol* **166**, 557-580.
- 5 Hansen, G. and Chilton, M. D. (1996). "Agrolistic" transformation of plant cells: integration of T-strands generated in planta. *Proc.Natl.Acad.Sci.U.S.A* **93**, 14978-14983.
- Hansen, G., Shillito, R. D., and Chilton, M. D. (1997). T-strand integration in maize protoplasts after codelivery of a T-DNA substrate and virulence genes.
- 10 *Proc.Natl.Acad.Sci.U.S.A* **94**, 11726-11730.
- Hanson, B., Engler, D., Moy, Y., Newman, B., Ralston, E., and Gutterson, N. (1999). A simple method to enrich an Agrobacterium-transformed population for plants containing only T-DNA sequences. *Plant J.* **19**, 727-734.
- Harlow, E. and Lane, D. (1988). "Antibodies: A Laboratory Manual." Cold Spring
- 15 Harbor Laboratory Press.
- Herrera-Estrella, L., De Block, M., Messens, E. H. J. P., Van Montagu, M., and Schell, J. (1983). Chimeric genes as dominant selectable markers in plant cells. *EMBO J.* **2**, 987-995.
- Hoffman, D. L., Laiter, S., Singh, R. K., Vaisman, I. I., and Tropsha, A. (1995).
- 20 Rapid protein structure classification using one-dimensional structure profiles on the bioSCAN parallel computer. *Comput.Appl.Biosci.* **11**, 675-679.
- Hooykens, P.J.J., Hall, M.A. & Libbeuga, K.R., eds. (1999) *Biochemistry and Molecular Biology of Plant Hormones* (Elsevier, Amsterdam).
- Houba-Heria, N., Pethe, C. d'Alayer, J & Lelouc, M. (1999) *Plant J.* **17**:615-626.
- 25 Klee, H.J. & Lanehon, M.B. (1995) in *Plant Hormones: Physiology, Biochemisry and Molecular Biology*, ed. Davies, P.J. (Kluwer, Dorddrocht, the Netherlands), pp. 340-353.

The first part of the report deals with the general situation of the country and the progress of the work during the year.

The second part of the report deals with the results of the work during the year.

The third part of the report deals with the conclusions of the work during the year.

- Krens, F. A., Molendijk, L., Wullems, G. J., and Schilperoort, R. A. (1982). *In vitro* transformation of plant protoplasts with Ti-plasmid DNA. *Nature* **296**, 72-74.
- 5 Lerner, R. A. (1982). Tapping the immunological repertoire to produce antibodies of predetermined specificity. *Nature* **299**, 593-596.
- Lerner, R. A., Green, N., Alexander, H., Liu, F. T., Sutcliffe, J. G., and Shinnick, T. M. (1981). Chemically synthesized peptides predicted from the nucleotide sequence of the hepatitis B virus genome elicit antibodies reactive with the native envelope protein of Dane particles. *Proc.Natl.Acad.Sci.U.S.A* **78**, 3403-3407.
- 10 Liddle, J. E. and Cryer, A. (1991). "A Practical Guide to Monoclonal Antibodies." Wiley New York.
- Loffler, J., Langui, D., Probst, A., and Huber, G. (1994). Accumulation of a 50 kDa N-terminal fragment of beta-APP695 in Alzheimer's disease hippocampus and neocortex. *Neurochem.Int.* **24**, 281-288.
- 15 Mok M.C. (1994) in *Cytokines: Chemistry, Activity and Function*, eds., Mok, D.W.S. & Mok,M.C. (CRC Boca Raton, Fl), pp.155-166.
- Monge, A., Lathrop, E. J., Gunn, J. R., Shenkin, P. S., and Friesner, R. A. (1995). Computer modeling of protein folding: conformational and energetic analysis of reduced and detailed protein models. *J.Mol.Biol* **247**, 995-1012.
- 20 Morris, R.O. et al. (1999). Isolation of a gene encoding a glycosylated cytokinin oxidase from maize. *Biochem. Biophys. Res. Commun.* **255**, 328-333
- Motyka, V., Faiss, M., Strnad, M., Kaminek, M. and Schmuelling, T. (1996). Changes in cytokinin content and cytokinin oxidase activity in response to derepression of *ipt* gene transcription in transgenic tobacco calli and plants. *Plant*
- 25 *Physiol.* **112**, 1035-1043.




- Murakami, T., Simonds, W. F., and Spiegel, A. M. (1992). Site-specific antibodies directed against G protein beta and gamma subunits: effects on alpha and beta gamma subunit interaction. *Biochemistry* **31**, 2905-2911.
- 5 Olszewski, K. A., Kolinski, A., and Skolnick, J. (1996). Folding simulations and computer redesign of protein A three-helix bundle motifs. *Proteins* **25**, 286-299.
- Osborne, B. I., Wirtz, U., and Baker, B. (1995). A system for insertional mutagenesis and chromosomal rearrangement using the Ds transposon and Cre-lox. *Plant J.* **7**, 687-701.
- 10 Ostresh, J. M., Blondelle, S. E., Dorner, B., and Houghten, R. A. (1996). Generation and use of nonsupport-bound peptide and peptidomimetic combinatorial libraries. *Methods Enzymol.* **267**, 220-234.
- Pabo, C. O. and Suchanek, E. G. (1986). Computer-aided model-building strategies for protein design. *Biochemistry* **25**, 5987-5991.
- 15 Paszkowski, J., Shillito, R. D., Saul, M., Mandak, V., and Hohn, T. H. B. P. I. (1984). Direct gene transfer to plants. *EMBO J.* **3**, 2717-2722.
- Peralta, E. G., Hellmiss, R., and Ream, W. (1986). Overdrive, a T-DNA transmission enhancer on the *A. tumefaciens* tumour-inducing plasmid. *EMBO J.* **5**, 1137-1142.
- 20 Quirino, B.F., Noh, Y.-S., Himmelbau, E., and Amasino, R.M. (2000). Molecular aspects of leaf senescence. *Trends in Plant Science* **5**, 278-282.
- Renouf, D. V. and Hounsell, E. F. (1995). Molecular modelling of glycoproteins by homology with non-glycosylated protein domains, computer simulated glycosylation and molecular dynamics. *Adv.Exp.Med.Biol* **376**, 37-45.
- 25 Rinaldi, A.C. and Comandini, O. (1999). Cytokinin oxidase strikes again. *Trends in Plant Sc.* **4**, 300.



- Rose, R. B., Craik, C. S., Douglas, N. L., and Stroud, R. M. (1996). Three-dimensional structures of HIV-1 and SIV protease product complexes. *Biochemistry* **35**, 12933-12944.
- 5 Rutenber, E. E., McPhee, F., Kaplan, A. P., Gallion, S. L., Hogan, J. C., Jr., Craik, C. S., and Stroud, R. M. (1996). A new class of HIV-1 protease inhibitor: the crystallographic structure, inhibition and chemical synthesis of an aminimide peptide isostere. *Bioorg.Med.Chem.* **4**, 1545-1558.
- Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). "Molecular Cloning: A Laboratory Manual." Cold Spring Harbor Laboratory Press.
- 10 Schlappi, M., Smith, D., and Fedoroff, N. (1993). TnpA trans-activates methylated maize Suppressor-mutator transposable elements in transgenic tobacco. *Genetics* **133**, 1009-1021.
- Shioda, T., Andriole, S., Yahata, T., and Isselbacher, K. J. (2000). A green fluorescent protein-reporter mammalian two-hybrid system with  
15 extrachromosomal maintenance of a prey expression plasmid: Application to interaction screening. *Proc.Natl.Acad.Sci.U.S.A* **97** , 5220-5224.
- Smulling, T., Rupp, H.M. Frank, M& Schafer, S.. (1999) in *Advances in Regulation of Plant Growth and Development*, eds. Surnad, M. Pac P. & Beck, E. (Peres, Prague), pp. 85-96.
- 20 Tamura, R. N., Cooper, H. M., Collo, G., and Quaranta, V. (1991). Cell type-specific integrin variants with alternative alpha chain cytoplasmic domains. *Proc.Natl.Acad.Sci.U.S.A* **88**, 10183-10187.
- Werner, T., Vadau Motyka, Miroslav Strnad, and Thomas Schmülling (2001) Regulation of plant growth by cytokinin. *Proc. Nat. Acad. Sci.*, **58 (18)** 10487-  
25 10492.
- Van Haaren, M. J., Sedee, N. J., Schilperoort, R. A., and Hooykaas, P. J. (1987). Overdrive is a T-region transfer enhancer which stimulates T-strand production in *Agrobacterium tumefaciens*. *Nucleic Acids Res.* **15**, 8983-8997.



- Van Sluys, M. A., Tempe, J., and Fedoroff, N. (1987). Studies on the introduction and mobility of the maize Activator element in *Arabidopsis thaliana* and *Daucus carota*. *EMBO J.* **6**, 3881-3889.
- 5 Wang, K., Genetello, C., Van Montagu, M., and Zambryski, P. C. (1987). Sequence context of the T-DNA border repeat element determines its relative activity during T-DNA transfer to plant cells. *Mol. Gen. Genet.* **210**, 338-346.
- 10 Woulfe, J., Lafortune, L., de Nadai, F., Kitabgi, P., and Beaudet, A. (1994). Post-translational processing of the neurotensin/neuromedin N precursor in the central nervous system of the rat—II. Immunohistochemical localization of maturation products. *Neuroscience* **60**, 167-181.
- Zhang, Y. L., Dawe, A. L., Jiang, Y., Becker, J. M., and Naider, F. (1996). A superactive peptidomimetic analog of a farnesylated dodecapeptide yeast pheromone. *Biochem. Biophys. Res. Commun.* **224**, 327-331.



**Annex to Form PCT/ISA/206  
COMMUNICATION RELATING TO THE RESULTS  
OF THE PARTIAL INTERNATIONAL SEARCH**

International Application No  
**PCT/EP 01/06833**

1. The present communication is an Annex to the invitation to pay additional fees (Form PCT/ISA/206). It shows the results of the international search established on the parts of the international application which relate to the invention first mentioned in claims Nos.:
- see 'Invitation to pay additional fees'
2. This communication is not the international search report which will be established according to Article 18 and Rule 43.
3. If the applicant does not pay any additional search fees, the information appearing in this communication will be considered as the result of the international search and will be included as such in the international search report.
4. If the applicant pays additional fees, the international search report will contain both the information appearing in this communication and the results of the international search on other parts of the international application for which such fees will have been paid.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 06571 A (UNIV MISSOURI) 11 February 1999 (1999-02-11) cited in the application	1, 2
Y	the whole document	7-10, 13-32, 34, 35, 39-62
X	ROY O. MORRIS ET AL.,: "Isolation of a gene encoding a glycosylated cytokinin oxidase from maize" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 255, 1999, pages 328-333, XP002151605 cited in the application	3-6, 11, 12
Y	* the whole document, in particular figure 6 *	7-10, 13-32, 34, 35, 39-62
X	DATABASE SWALL 'Online! 1 January 1998 (1998-01-01) S.D. ROUNSLEY ET AL.,: XP002151606 Database Accession number ID/AC=022213 abstract	3-6, 11, 12
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family



C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
✓ X	<p>DATABASE SWALL 'Online! 1 May 2000 (2000-05-01) M. BEVAN ET AL.,: "Cytokinin oxidase-like protein" XP002151607 Database Accession number ID/AC=Q9SU77 abstract</p>	3-6,11, 12
✓ X	<p>DATABASE SWALL 'Online! 1 May 1999 (1999-05-01) X. LIN ET AL.: "Arabidopsis thaliana chromosome II BAC F3P11 genomic sequence. Putative Cytokinin Oxidase" XP002151608 Database Accession number ID/AC=Q9ZUP1 abstract</p>	3-6,11, 12
✓ X	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; M. FRANK ET AL.,: "TSD genes negatively regulate meristematic activity in Arabidopsis" XP002151616 Database Accession number AN=PREV200000242628 * abstract * &amp; Biologia Plantarum (Prague) 1999, Vol. 42 (Suppl.), page S47</p>	11,12
✓ A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; ZHANG, N. ET AL.,: "Initiation and elongation of lateral roots in Lactuca sativa" XP002151609 Database Accession number AN=PREV199900326622 * abstract * &amp; International J. of Plant Sciences 1999, Vol. 160(3), pages 511-519</p> <p style="text-align: center;">-/-</p>	1,2, 15-30, 39-62



Annex to Form PCT/ISA/206  
COMMUNICATION RELATING TO THE RESULTS  
OF THE PARTIAL INTERNATIONAL SEARCH

International Application No

PCT/EP 01/06833

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; Y. KODA ET AL.,: "Cytokinin production by tomato root. Identification of a major cytokinin produced by the root and environmental factors affecting the production" XP002151610 Database Accession number AN=PREV198988038194 * abstract * &amp; J. of the Faculty of Agriculture Hokkaido University 1989, Vol. 64(1), pages 10-20</p> <p>-----</p>	<p>1,2, 15-30, 39-62</p>



**Patent Family Annex**  
Information on patent family members

International Application No  
**PCT/EP 01/06833**

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9906571	A	11-02-1999	US 6229066 B1	08-05-2001
			AU 8603898 A	22-02-1999
			BR 9811592 A	19-09-2000
			CN 1265146 T	30-08-2000
			EP 1002096 A1	24-05-2000
			PL 338330 A1	23-10-2000
			WO 9906571 A1	11-02-1999
			HU 0002349 A2	28-11-2000

